CUMULATIVE AUTHOR INDEX
Environmental Entomology
Volume 41, 2012

A

Abell, K. J., 925
Abrahamson, M. D., 750
Accamando, A. K., 551
Adamo, S. A., 1644
Adamson, N. L., 813
Addesso, K. M., 887
Alatorre-Rosas, R., 959
Alcantara, E. P., 1268
Aldana, A., 497
Allan, B. F., 1062
Allen, K. C., 1302
Allison, J. D., 1587
Altuntaş, H., 688
Álvarez-Alfageme, F., 1687
Amarasekare, K. G., 1184
Anderson, F. E., 723
Andjelkovic, M., 979
Andow, D. A., 1045
Arbogast, R. T., 152
Arioli, C. J., 849
Arzuffi, R., 387
Asokan, R., 1239
Aukema, B. H., 578

B

Ba, N. M., 1255
Bacher, S., 1086
Backus, E. A., 1215
Badenhausser, I., 1050
Bae, Y. J., 950
Baek, M. J., 950
Bahar, M. H., 1653
Baker, J. L., 1644
Ballman, E. S., 1417
Barman, A. K., 125
Barrett, B. A., 933
Barzegar, M., 1638
Bauce, E., 1290
Beaudoin, A.L.P., 362
Beauzay, P. B., 282
Beck, J. J., 603
Beeman, R. W., 188
Beers, E. H., 1364
Beizhou, S., 872
Bejleri, K., 802
Bennett, A. B., 1077
Bento, J.M.S., 562
Berenbaum, M. R., 1435
Berg, S. H., 1466
Bergmann, D. J., 1069
Berkov, A., 108
Berthiaume, R., 1290
Bextine, B., 98
Bhat, P. S., 1239
Biazzo, J., 665
Bickerton, M. W., 612
Bigler, F., 1687
Bisognin, M., 849
Blais, J., 1290
Blankenship, E. E., 1494
Blossey, B., 1086
Boiteau, G., 916
Boivin, G., 159
Bonte, J., 989
Boone, J., 594
Booth, W., 426
Bordera, S., 587
Botton, M., 849
Boughton, A. J., 308
Bourgeois, G., 159
Bradford, M. A., 463
Bradshaw, J. D., 865
Braman, S. K., 1145, 1153
Brandenburg, D., 1069
Brandsch, C., 676
Bretagnolle, V., 1050
Brodeur, J., 159, 1290
Buckman, K., 750
Burbano, E. G., 133
Burkness, E. C., 172
Burrack, H. J., 265
Buss, L. J., 487

C

Cabezas, M. F., 849
Cáceres, C. E., 497
Calderon, O., 108
Campbell, J. F., 188, 1501
Capinera, J. L., 1131
Carleton, D., 594
Carlson, B., 180
Carroll, A. L., 392
Carvalho, V.F.P., 1454
Casagrande, R. A., 841
Casteel, C. L., 1435
Castrillo, L. A., 822
Cervenka, V. J., 91
Chavigny, P., 420
Cheah, C., 1662
Chebny, V., 399
Chen, H., 697
Chen, J., 1008
Chen, M., 880
Chen, P., 1606
Chen, S.-J., 469
Chen, X. M., 1163
Cheng, J.-A., 469
Cherry, R., 887
Chi, H., 30
Choate, B., 222
Clark, E. L., 392, 1386
Clark, R. E., 1405
Clayton, G. W., 72
Clercq, P. D., 989
Cockfield, S. D., 1364
Cognato, A. I., 776
Collins, H. L., 880
Cong, L., 1672
Conlong, D., 989
Contreras-Garduño, J., 959
Cooper, W. R., 342, 941
Córdoba-Aguilar, A., 959
Cornulier, T., 1050
Corsato, C.D.A., 233
Cossé, A. A., 648
Coyle, D. R., 91, 731
Cronin, J. T., 551
Crook, D. J., 648, 1311
Crosslin, J. M., 1019
Cuddington, K., 508, 516
Cui, H., 1443
Cutler, G. C., 370, 1097

D

Daane, K. M., 81, 497
Dabire, C., 1255
D'Amico, V., 1466
Daniell, T. J., 1386
Davis, C., 289
De Bortoli, C. P., 1454
De Bortoli, S. A., 1454
Delfín-González, H., 587
Delucia, E. H., 1435
De Souza, A. R., 233
Deyrup, M. A., 1597
Dhileepan, K., 317
Doak, P., 180
Dodds, K. J., 1350
Dosdall, L. M., 72, 1653
Dowd, S. E., 98
Dowling, L., 1169
Drabble, S., 289
Drees, B. M., 1008
Dreves, A. J., 621
Driesche, R.G.V., 925
Drummond, F. A., 59, 222
Duan, J. J., 245, 925
Dubois, G. D., 1350
Dudley, N., 133
Duehl, A., 152

E

Earle, E. D., 880
El-Shazly, M. M., 11
Emerson, S. C., 355
Encalada, A. C., 1322

CUMULATIVE AUTHOR INDEX
Environmental Entomology
Volume 41, 2012

A

Abell, K. J., 925
Abrahamson, M. D., 750
Accamando, A. K., 551
Adamo, S. A., 1644
Adamson, N. L., 813
Addesso, K. M., 887
Alatorre-Rosas, R., 959
Alcantara, E. P., 1268
Aldana, A., 497
Allan, B. F., 1062
Allen, K. C., 1302
Allison, J. D., 1587
Altuntaş, H., 688
Álvarez-Alfageme, F., 1687
Amarasekare, K. G., 1184
Anderson, F. E., 723
Andjelkovic, M., 979
Andow, D. A., 1045
Arbogast, R. T., 152
Arioli, C. J., 849
Arzuffi, R., 387
Asokan, R., 1239
Aukema, B. H., 578

B

Ba, N. M., 1255
Bacher, S., 1086
Backus, E. A., 1215
Badenhausser, I., 1050
Bae, Y. J., 950
Baek, M. J., 950
Bahar, M. H., 1653
Baker, J. L., 1644
Ballman, E. S., 1417
Barman, A. K., 125
Barrett, B. A., 933
Barzegar, M., 1638
Bauce, E., 1290
Beaudoin, A.L.P., 362
Beauzay, P. B., 282
Beck, J. J., 603
Beeman, R. W., 188
Beers, E. H., 1364
Beizhou, S., 872
Bejleri, K., 802
Bennett, A. B., 1077
Bento, J.M.S., 562
Berenbaum, M. R., 1435
Berg, S. H., 1466
Bergmann, D. J., 1069
Berkov, A., 108
Berthiaume, R., 1290
Bextine, B., 98
Bhat, P. S., 1239
Biazzo, J., 665
Bickerton, M. W., 612
Bigler, F., 1687
Bisognin, M., 849
Blais, J., 1290
Blankenship, E. E., 1494
Blossey, B., 1086
Boiteau, G., 916
Boivin, G., 159
Bonte, J., 989
Boone, J., 594
Booth, W., 426
Bordera, S., 587
Botton, M., 849
Boughton, A. J., 308
Bourgeois, G., 159
Bradford, M. A., 463
Bradshaw, J. D., 865
Braman, S. K., 1145, 1153
Brandenburg, D., 1069
Brandsch, C., 676
Bretagnolle, V., 1050
Brodeur, J., 159, 1290
Buckman, K., 750
Burbano, E. G., 133
Burkness, E. C., 172
Burrack, H. J., 265
Buss, L. J., 487

C

Cabezas, M. F., 849
Cáceres, C. E., 497
Calderon, O., 108
Campbell, J. F., 188, 1501
Capinera, J. L., 1131
Carleton, D., 594
Carlson, B., 180
Carroll, A. L., 392
Carvalho, V.F.P., 1454
Casagrande, R. A., 841
Casteel, C. L., 1435
Castrillo, L. A., 822
Cervenka, V. J., 91
Chavigny, P., 420
Cheah, C., 1662
Chebny, V., 399
Chen, H., 697
Chen, J., 1008
Chen, M., 880
Chen, P., 1606
Chen, S.-J., 469
Chen, X. M., 1163
Cheng, J.-A., 469
Cherry, R., 887
Chi, H., 30
Choate, B., 222
Clark, E. L., 392, 1386
Clark, R. E., 1405
Clayton, G. W., 72
Clercq, P. D., 989
Cockfield, S. D., 1364
Cognato, A. I., 776
Collins, H. L., 880
Cong, L., 1672
Conlong, D., 989
Contreras-Garduño, J., 959
Cooper, W. R., 342, 941
Córdoba-Aguilar, A., 959
Cornulier, T., 1050
Corsato, C.D.A., 233
Cossé, A. A., 648
Coyle, D. R., 91, 731
Cronin, J. T., 551
Crook, D. J., 648, 1311
Crosslin, J. M., 1019
Cuddington, K., 508, 516
Cui, H., 1443
Cutler, G. C., 370, 1097

D

Daane, K. M., 81, 497
Dabire, C., 1255
D'Amico, V., 1466
Daniell, T. J., 1386
Davis, C., 289
De Bortoli, C. P., 1454
De Bortoli, S. A., 1454
Delfín-González, H., 587
Delucia, E. H., 1435
De Souza, A. R., 233
Deyrup, M. A., 1597
Dhileepan, K., 317
Doak, P., 180
Dodds, K. J., 1350
Dosdall, L. M., 72, 1653
Dowd, S. E., 98
Dowling, L., 1169
Drabble, S., 289
Drees, B. M., 1008
Dreves, A. J., 621
Driesche, R.G.V., 925
Drummond, F. A., 59, 222
Duan, J. J., 245, 925
Dubois, G. D., 1350
Dudley, N., 133
Duehl, A., 152

E

Earle, E. D., 880
El-Shazly, M. M., 11
Emerson, S. C., 355
Encalada, A. C., 1322

Enríquez-Vara, J. N., 959
Epsky, N. D., 1184, 1597
Erbilgin, N., 478, 1575
Ergin, E., 688
Errouissi, F., 785
Espírito-Santo, M. M., 541
Estes, A. M., 265

F

Fell, R. D., 813
Feng, H., 1606
Feng, J.-N., 706
Fernandes, F. L., 333
Fernandes, M.E.S., 333
Fierke, M. K., 1311
Fisher, G. C., 621
Flinn, P. W., 1501
Fliszkiewicz, M., 1621
Fombong, A. T., 152
Frank, S. D., 377
Freier, B., 676
Frost, K. E., 1553
Fuester, R., 245

G

Gabel, M., 1069
Galligan, L. D., 1311
Gandhi, K.J.K., 731
Gao, L., 349
Garcia, M. S., 849
García-López, A., 1544
Gardner, W. A., 298
Garnas, J., 59
Garner, S. R., 1680
Garrido-Jurado, I., 966
Ge, F., 714, 1029, 1443
Gent, D. H., 621
Gerber, E., 1086
Ghising, K., 282, 1417
Giaccai, J., 1409
Giejdasz, K., 1621
Giladi, I., 463
Gilg, M. R., 1246
Gillette, N. E., 133, 1575
Giustolin, T. A., 233
Glemser, E. J., 1169
Goarant, A., 1050
Gonda-King, L., 523
Gontijo, L. M., 1364
González-Moreno, A., 587
Goodisman, M.A.D., 1612
Goodsman, D. W., 478
Gouat, M., 1050
Goyal, G., 1131
Gratton, C., 1077
Gray, M. E., 865
Grevstad, F. S., 448
Griggs, M. H., 822
Groden, E., 59
Groot, P. D., 289
Groves, R. L., 532, 1215, 1279, 1553
Guangbo, T., 872
Guillén, L., 1597
Guo, J.-Y., 1672
Guo, X., 1606
Guzmán-Franco, A. W., 959

H

Haavik, L. J., 1311
Hagler, J. R., 1279, 1527
Hahn, N. G., 455
Hail, D., 98
Hajek, A. E., 1115
Hale, R. J., 1190
Hallett, R. H., 1169
Hamilton, G. C., 612
Hamm, J. C., 571
Hanks, L. M., 1587
Hanula, J. L., 40
Harbur, M. M., 750
Harker, K. N., 72
Harmon, J. P., 282, 1417, 1426
Hay-Roe, M. M., 487
Hazlehurst, A. F., 841
Hearn, D. J., 265
Hébert, C., 1290
Heimpel, G. E., 750
Hein, G. L., 1494
Held, D. W., 1139
Hellmich, R. L., 200, 1047
Helm, S., 1462
Helms, T. C., 282
Henneken, R., 1462
Herbert, D. A., 761
Herms, D. A., 1372
Herrick, N. J., 118
Higgs, K. M., 355
Hinz, H. L., 1086
Hoback, W. W., 238
Hoebeke, E. R., 1350
Holmes, L. A., 971
Hong, X.-Y., 833
Horn, S., 40
Hou, M., 1231
Hough-Goldstein, J. A., 1466
Hu, C., 1231
Hu, X. P., 1139
Huang, Z. Y., 1163
Hubbard, S. F., 1386
Huber, D.P.W., 392
Hullé, M., 1398
Hummel, J. D., 72
Hummel, N. A., 571
Hunt, T. E., 1494
Huseth, A. S., 532, 1553
Hutchison, W. D., 172, 200

I

Inglis, D., 1169
Inouye, D. W., 1332
Isaacs, R., 455

J

Jackai, L., 1255
Jackson, R. E., 1302
James, D. G., 621
Jensen, A. S., 768
Jha, R. K., 30
Jiang, H.-X., 706
Jiang, M.-X., 469
Jiang, X., 1199
Jie, Z., 872
Jin, Q.-A., 1208
Jin, T., 1208
Johns, R. C., 594
Johnson, M. W., 497, 1215, 1279
Jones, T., 133
Jones, W. A., 1008
Journey, A. M., 91

K

Kajimura, H., 255
Kang, J., 200, 1255
Kanzaki, N., 828
Kaplan, I., 1527
Karley, A. J., 1386
Katz, M. L., 448
Kaufmann, M., 133
Keena, M. A., 1662
Keesey, I. W., 933
Kelley, S. T., 723
Kelly, J. L., 1527
Kendra, P. E., 1597
Kenig, B., 979
Kennedy, G. G., 362
Kevan, P., 855
Khosla, S., 855
Khrimian, A., 648
Kiliç, A. Y., 688
King, J. R., 1405
Kiowsi, A., 676
Klose, T., 676
Knauft, D. A., 1145, 1153
Knight, A. L., 399, 407
Knodel, J. J., 282
Knuteson, D. L., 1553
Koch, R. L., 91, 750
Koenig, R. L., 487
Kok, L. T., 118, 896
Kovacs, J. L., 1612
Krengel, S., 676
Krishnan, N., 1621
Krugner, R., 1279
Kuhar, T. P., 761
Kumar, N.K.K., 1239

L

Labidi, I., 785
Lake, E., 1466
Lamp, W. O., 802
Leach, E. L., 1680
Lee, C.-Y., 426
Legaspi, Jr., B. C., 669
Legault, S., 1290
Leggo, J. J., 594

Lei, X., 349
Leirana-Alcocer, J., 587
Lepage, M. P., 159
Lerch, R. N., 933
Leslie, A. W., 802
Levy, K., 81
Li, B.-L., 714
Li, C., 1443
Li, D., 349
Li, G., 1606
Li, K., 1163
Li, X.-W., 706
Li, Z., 349
Lieffers, V. J., 478
Light, D. M., 399, 407, 603
Lin, C.-H., 933
Lin, Y.-Y., 1208
Lind, E. M., 1409
Lindgren, B. S., 578
Liu, X., 880
Long, Y., 1231
Lopes, E. N., 233
Lopes-Mielezrski, G. N., 233
Lorenzen, M. D., 188
Lövei, G. L., 1045
Lovett, M.M.E., 1644
Lowenstein, D. M., 532
Lu, F., 469
Ludoski, J., 979
Luo, C. W., 1163
Luo, L., 1199
Luo, S.-P., 657
Luttrell, R. G., 1302
Luvizotto, R. A., 562
Lynch, D. H., 370, 1097

M

Machial, L. A., 578
MacKenzie, K., 370, 1097
MacKinley, P., 916
Makni, H., 420
Makni, M., 420
Mannion, C. M., 1184
Marcos-García, M. Á., 1544
Martinez, H.E.P., 333
Masuya, H., 255
Mayo, P., 648
Mcauslane, H. J., 887
McAvoy, T. J., 118
McClay, A. S., 317
McCullough, D. G., 1115
McElfresh, J. S., 1587
McEvoy, P. B., 355
McFadden-Smith, W., 1169
McFadyen, R. E., 317
McGarvey, B., 855
McGrath, J. M., 802
McKenney, J. L., 1587
McPhee, K., 59
Meagher, R. L., 1131
Medina, R. F., 125
Mehmel, C. J., 1575
Mehrkhou, F., 326
Meissle, M., 1037
Melo, G. A., 541
Menzel, A., 1462
Merrill, S. C., 1505
Milbrath, L. R., 665
Millar, J. G., 1587
Miller, N. J., 997
Mills, N. J., 81
Mitchell, R. F., 1587
Moharramipour, S., 326, 1638
Moll, E., 676
Montgomery, M. E., 1662
Montgomery, W. S., 1597
Morey, A. C., 172
Mori, S. R., 133, 1575
Morse, A., 855
Morse, J. G., 51, 1279
Moser, S. E., 200
Muilenburg, V. L., 1372
Mullins, D. E., 813
Munyaneza, J. E., 1019
Murphy, A. F., 768
Murtaugh, P. A., 355
Muturi, E. J., 1062
Myron, E. P., 1409

N

Nachman, G., 20, 141
Nair, S., 1145, 1153
Naranjo, S. E., 1047
Nardi, C., 562
Nava, D. E., 849
Naveh, V. H., 326
Needelman, B. A., 802
Niogret, J., 1597
Niziolek, O. K., 1435
Nott, R. W., 289
Nouira, S., 785
Novicic, Z. K., 979
Nuessly, G. S., 1131

O

O'Casey, C., 448
Ochoa, I., 289
Ode, P. J., 282
O'Donovan, J. T., 72
Oi, D. H., 1008
Oliveira, K. N., 541
Oliver, J. B., 636
O'Neal, M. E., 1565
Onstad, D. W., 200, 880, 1255
Othman, A. S., 426
Ouyang, F., 1029
Owen, D. R., 1575

P

Pahs, T. L., 91
Pálinkás, Z., 1687
Parajulee, M. N., 125, 1029
Parker, J. D., 1409
Parra, J.R.P., 562
Paula-Moraes, S. V., 1494
Peairs, F. B., 1505
Pemberton, R. W., 308
Peng, Z.-Q., 1208
Perera, O. P., 1302
Permkam, S., 776
Petit, S., 1069
Pfammatter, J. A., 91
Philips, C. R., 761
Philpott, S. M., 1107
Picanço, M. C., 333
Pickering, G. J., 1169
Pierson, E. A., 265
Pinto-Leite, C. M., 1534
Pittendrigh, B. R., 1255
Pomari, A. F., 1454
Popay, A., 433
Porter, P. M., 750
Prasifka, J. R., 200, 865
Prat, N., 1322
Preisser, E. L., 523
Prischmann-Voldseth, D. A., 282, 1417
Pyke, G. H., 1332

Q

Qiu, B., 657
Qiu, F., 1606
Queiroz, R. B., 333
Querino, R. B., 233
Quesada-Moraga, E., 966
Quiring, D. T., 594

R

Raboudi, F., 420
Radville, L., 523
Ragsdale, D. W., 750
Ramalho, D. G., 1454
Ramamurthy, V. V., 1239
Raman, A., 433
Ranger, C. M., 636
Rao, S., 905
Reagan, T. E., 571
Reay-Jones, F.P.F., 1516
Rebijith, K. B., 1239
Reding, M. E., 636
Reese, K. M., 1107
Reeve, J. D., 723
Reisig, D. D., 761
Reitz, S. R., 487
Rempoulakis, P., 265
Rendón, P. A., 497
Renkema, J. M., 370, 1097
Reynolds, P. G., 508, 516
Ricci, J., 1062
Ríos-Touma, B., 1322
Ro, M. D., 989
Roberts, E. A., 761
Robledo, N., 387
Rocha, P.L.B., 1534
Romeis, J., 1037, 1047, 1687
Rondon, S. I., 768

Rossi, A. M., 1246
Rotheray, G. E., 1544
Roulston, T. H., 813
Roush, R., 880
Ruppert, D. E., 802
Ryall, K. L., 648
Ryan, K., 289

S

Saeidi, F., 1638
Sahashi, N., 828
Salom, S. M., 118, 896
Sansone, C. G., 125
Sappington, T. W., 997, 1199
Scaduto, D. A., 1680
Scarr, T., 648
Schultz, P. B., 636
Seal, D. R., 1131
Sears, M. K., 1169
Segal, L. M., 1435
Semeao, A. A., 188
Senaratne, K.A.D.W., 317
Sezen, Z., 750
Sformo, T., 180
Shapiro, J. P., 487
Shapiro-Ilan, D. I., 298
Shelton, A. M., 880, 1047
Shen, Z., 349
Shi, B., 1231
Shi, P., 714
Shipp, L., 855
Shirk, P. D., 487
Sidhu, J. K., 571
Siegert, N. W., 1115
Silk, P. J., 648
Silva, J. O., 541
Simmons, A. M., 669
Singham, G. V., 426
Sisterson, M. S., 1279
Sittichaya, W., 776
Skovgård, H., 20, 141
Sloderbeck, P. E., 188
Smith, R. F., 802
Smith, S. M., 289, 594
Snodgrass, G. L., 1302
Snyder, A. L., 118
Soliman, M. M., 11
Son, Y., 1215
Soroka, J. J., 1653
Spurgeon, D. W., 342, 415, 941
Srikumar, K. K., 1239
Srygley, R. B., 166
Stamenkovic-Radak, M., 979
Stangl, G. I., 676
Steck, G. J., 1131
Steenweg, R. W., 578
Steiner, P. M., 180
Stephen, F. M., 1311
Stiling, P., 1246
Stokes, K., 1246
Stork, R., 1631
Story, H. M., 896
Stout, M. J., 571
Strange, J. P., 905
Su, J., 1443
Sun, J., 997
Sun, Y., 1443
Sweeney, J., 648
Swisher, K. D., 1019

T

Talebi, A. A., 326
Tamò, M., 1255
Tanaka, R., 828
Tang, L.-C., 30
Taylor, D. B., 213
Teal, P.E.A., 152
Teulon, D.A.J., 1190
Tewksbury, L., 841
Thomas, J.M.G., 487
Thompson, G. J., 1680
Thomson, J. D., 1332
Throne, J. E., 697
Tomberlin, J. K., 971
Tomisawa, N., 255
Torto, B., 152
Tran, L. T., 1190
Trevino, M., 317
Trimble, R. M., 1481
Trotter, III, R. T., 1662
Turgeon, J. J., 289

U

Uçkan, F., 688
Ugine, T. A., 1
Uma, D. B., 1474

V

Vacari, A. M., 1454
Valverde-García, P., 966
Vandenberg, J. D., 822
Vanlaerhoven, S. L., 971
Vanlerberghe, F. M., 420
Varenhorst, A. J., 1565
Vargo, E. L., 426
Vasquez, C. J., 51
Venette, R. C., 172
Vieira, L. C., 896
Vitelli, M. P., 317

W

Wagner, D., 180
Walde, S. J., 370, 1097
Wan, F.-H., 1125, 1672
Wang, X.-G., 81, 497
Warren, II, R. J., 463
Wasielewski, O., 1621
Watt, T., 925
Way, M. O., 469
Webster, J. N., 1575
Weed, A. S., 841
Weiss, M. R., 1474
Wells, L., 298
Wen, H.-B., 1208
Whalen, R., 1426
Wheatley, W., 433
Wheeler, M. M., 1115
Whipple, S. D., 238
White, W. H., 275
Whitworth, R. J., 188
Wienhold. B. J., 213
Wiggins, N. L., 872
Willms, S., 152
Wilson, G., 1644
Wilson, L. T., 275
Wishart, J., 1386
Wood, B. W., 298
Wood, D. L., 1575
Woods, J. L., 621
Worner, S. P., 1190
Wright, L. C., 621
Wright, M. G., 133
Wright, R. J., 1494
Wyckhuys, K.A.G., 750
Wyman, J. A., 1553

X

Xavier, V. M., 333
Xie, D., 1199
Xu, Y., 1139
Xu, Z.-F., 657
Xusheng, S., 872

Y

Yamaoka, Y., 255
Yang, B., 1029
Yang, S., 925
Yang, X., 1231
Yokoyama, V. Y., 497, 1177
Yokoyama-Hatch, H. A., 497
Yoon, T. J., 950
Yuncong, Y., 872
Yurchenko, G., 245

Z

Zalom, F. G., 349
Zanardi, O. Z., 849
Zayed, A., 11
Zhang, G.-F., 1125
Zhang, H., 697
Zhang, L., 1199
Zhang, X.-C., 706
Zhang, X.-F., 833
Zhang, Y., 40
Zhao, D.-X., 833
Zhao, L., 1008
Zhou, Z.-S., 657, 1672
Zhu, K. Y., 697
Zucchi, R. A., 233
Zumbado, M. A., 1544
Zylstra, K. E., 1350

CUMULATIVE KEY WORD INDEX
Environmental Entomology
Volume 41, 2012

A

abiotic factors,
dung beetle assemblages, 785
Mediterranean bioclimate, 785
spatial scale, 785
temporal scale, 785
activity,
Braconidae, 587
Ichneumonidae, 587
relative humidity, 587
temperature, 587
Adelges tsugae,
false rings, 523
Fiorinia externa, 523
herbivory, 523
Tsuga canadensis, 523
Adelphocoris suturalis,
Apolygus lucorum, 1606
diapause, 1606
photoperiod, 1606
sensitive stage, 1606
adult emergence,
egg clustering, 971
egg desiccation hypothesis, 971
egg eclosion, 971
waste management, 971
adult trees,
environmental condition, 828
pine wood nematode, 828
shading, 828
virulence, 828
Aethina tumida,
honey bees, 152
insect trapping, 152
small hive beetle larvae, 152
Aganaspis,
Anastrepha, 233
Asobara, 233
Doryctobracon, 233
Neosilba, 233
age-specific life table,
development, 1
fecundity, 1
Lygus lineolaris, 1
temperature, 1
aggregation,
Blissus insularis Barber, 887
host-plant volatiles, 887
pheromones, 887
Y-tube olfactometer, 887
aggressive mimicry,
anti-predator behavior, 1474
mud-dauber wasps, 1474
predator–prey interaction, 1474
Theridiidae, 1474
agricultural areas,
diversity, 776
flight pattern, 776
Xyleborini ambrosia beetles, 776
agriculture,
Diuraphis noxia, 1505
pest, 1505
Russian wheat aphid, 1505
spatial model, 1505
Agrilus planipennis, 245, 925
biological control, 925
exotics and invasive species, 245
indigenous, 245
interspecific competition, 925
parasitoids, 245
Spathius galinae, 925
Tetrastichus planipennisi, 925
wood borers, 245
agroforestry,
Azteca instabilis, 1107
coffee, 1107
Pseudacteon, 1107
trait-meditated interactions, 1107
Ailanthus altissima,
classical biological control, 118
Eucryptorrhynchus brandti, 118
host-specificity, 118
invasive species, 118
alkaloids,
coccid, 333
coffee plants, 333
nutrients, 333
phenolic compounds, 333
ambrosia beetle, 133, 636, 822
artificial diet, 822
ethanol, 636
invasive pest, 822
laboratory rearing, 822
semiochemicals, 133
solid phase microextraction, 636
traps, 133
Xylosandrus germanus, 636
Anastrepha,
Aganaspis, 233
Asobara, 233
Doryctobracon, 233
Neosilba, 233
ancestral state reconstruction,
bark beetles, 723
conifers, 723
Dendroctonus, 723
outbreaks, 723
ant diversity,
Formicidae, 222
lowbush blueberry, 222
Maine, 222
antibiotics,
Lissorhoptrus oryzophilus, 469
reproduction, 469
symbiont, 469
Wolbachia, 469
ant–plant interactions,
elaiosome, 1405
mutualism, 1405
myrmecochory, 1405
trophic interactions, 1405
anti-predator behavior,
aggressive mimicry, 1474
mud-dauber wasps, 1474
predator–prey interaction, 1474
Theridiidae, 1474
Aonidiella aurantii,
biological control, 51
longevity, 51
sex ratio, 51
size, 51
Aphaenogaster picea,
Aphaenogaster rudis, 463
biotic interactions, 463
community ecology, 463
Lasius alienus, 463
Aphaenogaster rudis,
Aphaenogaster picea, 463
biotic interactions, 463
community ecology, 463
Lasius alienus, 463
aphid, 59, 1077, 1638
flower diversity, 1077
Homoptera, 59
invasive ant, 59
mannitol, 1638
myrmecophily, 59
Myrmica rubra, 59
natural enemy diversity, 1077
natural pest suppression, 1077
overwintering, 1638
polyols, 1638
supercooling points, 1638
Aphis glycines, 282, 750, 1417, 1426
Binodoxys communis, 282, 1417
biological control, 282, 1417
Cerotoma trifurcata, 750
Empoasca fabae, 750
host-plant resistance, 282, 1417, 1426

movement, 1426
soybean, 282, 750, 1417
winter rye, 750
within-plant distribution, 1426
Apis mellifera,
distribution, 1462
phenology, 1462
swarm, 1462
Apolygus lucorum,
Adelphocoris suturalis, 1606
diapause, 1606
photoperiod, 1606
sensitive stage, 1606
apoptosis,
encapsulation, 688
Galleria mellonella, 688
gibberellic acid, 688
hemocyte, 688
apparent mortality,
population dynamics, 1311
Quercus rubra, 1311
scar tissue, 1311
wood borers, 1311
apple, 399, 407, 1364
biological control, 1364
Cydia pomonella, 399, 407
Eriosoma lanigerum, 1364
mating disruption, 399, 407
monitoring, 399
natural enemies, 1364
plant volatiles, 407
sex pheromone, 399
survey, 1364
traps, 407
apple orchard,
aromatic plant, 872
arthropod community, 872
herbivore, 872
predator, 872
Araneae,
Atlantic Rainforest, 1534
day collections, 1534
night collections, 1534
sampling effort, 1534
aromatic plant,
apple orchard, 872
arthropod community, 872
herbivore, 872
predator, 872
arrestment,
kairomone, 603
larvae, 603
microencapsulated, 603
pear ester, 603
arthropod abundance,
Bt corn, 1268
monitoring, 1268
PRC analysis, 1268
riparian area, 1268
arthropod community,
apple orchard, 872
aromatic plant, 872
herbivore, 872
predator, 872
artificial diet, 30, 265, 822
ambrosia beetle, 822
Helicoverpa armigera, 30
invasion ecology, 265
invasive pest, 822
laboratory rearing, 822
life table, 30
olive fruit fly, 265
sterile insect technology, 265
symbiosis, 265
Zea mays, 30
Asobara,
Aganaspis, 233
Anastrepha, 233
Doryctobracon, 233
Neosilba, 233
Asopinae,
biological control, 1454
Brassicaceae, 1454
aspen leaf miner,
microhabitat preference, 180
overwintering, 180
Populus tremuloides, 180
supercooling, 180
Asphondylia borrichiae,
host-associated divergence, 1246
host-race formation, 1246
sympatric speciation, 1246
Atlantic Rainforest,
Araneae, 1534
day collections, 1534
night collections, 1534
sampling effort, 1534
augmentation,
biological control, 1527
ELISA, 1527
predation monitoring, 1527
protein marking, 1527
Aulacorthum solani,
foxglove aphid, 665
host suitability, 665
swallow-wort, 665
Vincetoxicum, 665
autecology,
habitat, 966
integrated pest management (IPM), 966
soil-dwelling pest, 966
Azteca instabilis,
agroforestry, 1107
coffee, 1107
Pseudacteon, 1107
trait-meditated interactions, 1107

B

Bacillus thuringiensis,
Cry3Bb1, 1037
nontarget risk assessment, 1037
Theridion impressum, 1037
bacterial community,
Brevicoryne brassicae, 1386
endosymbiont, 1386
real-time qPCR, 1386
16S ribosomal DNA, 1386
Bactericera cockerelli, 1019, 1190
development, 1190
liberibacter, 1019
potato, 1019
psyllid haplotypes, 1019
temperature, 1190
tomato psyllid (potato psyllid), 1190
zebra chip, 1019
Bactrocera oleae, 497, 1177
cultural control, 1177
fruit fly dispersal, 1177
irradiated host, 497
larval parasitoid, 497
longevity, 497
mature larvae, 1177
parasitism, 497
teneral adult, 1177
Bailey's triple-catch, 20, 141
cattle, 20
fluorescent marking, 20
forecasting, 20
Jolly-Seber's stochastic method, 141
longevity, 20
loss rate, 141
Monte-Carlo simulation, 141
population estimates, 141
banker plants,
Frankliniella, 487
minute pirate bugs, 487
Orius, 487
thrips, 487
barcode,
mycetome, 108
Peribocum pubescens, 108
symbiont, 108
Xylergates pulcher, 108
bark beetles,
ancestral state reconstruction, 723
conifers, 723
Dendroctonus, 723
outbreaks, 723
bean plataspid,
exotic, 40
host preference, 40
invasive, 40
kudzu bug, 40

Beauveria bassiana,
biological control, 298
cover crop, 298
Curculio caryae, 298
pecan, 298
beet armyworm,
Iran, 326
life table, 326
reproduction, 326
soybean, 326
beet leafhopper,
Circulifer, 768
nonparametric multiplicative regression, 768
pest management, 768
purple top disease, 768
beetle-fungi association,
Fusarium solani, 255
phoretic, 255
Scolytogenes birosimensis, 255
tree mortality, 255
Bemisia tabaci, 669, 1443
biological control, 669
Delphastus catalinae, 669
diet, 669
jasmonic acid, 1443
predator, 669
salicylic acid, 1443
tomato, 1443
whitefly preconditioning, 1443
Bemisia tabaci biotype B, 1125, 1672
high-temperature stress, 1672
host suitability, 1125
interspecific interaction, 1125
leaf age, 1125
life table, 1672
long-term fitness effects, 1672
Trialeurodes vaporariorum, 1125
bermudagrass,
Eisenia fetida, 1139
feeding ecology, 1139
plant insect interactions, 1139
Scapteriscus, 1139
Betula,
forest decline, 1372
host plant resistance, 1372
insect outbreaks, 1372
population dynamics, 1372
Binodoxys communis, 282, 1417
Aphis glycines, 282, 1417
biological control, 282, 1417
host-plant resistance, 282, 1417
soybean, 282, 1417
biocontrol,
dispersal, 1466
orientation, 1466
Persicaria perfoliata, 1466
Rhinoncomimus latipes, 1466
biodiversity,
drainage ditch, 802
macroinvertebrate, 802
nutrients, 802
water quality, 802
bioenergy,
ethanol, 865
herbivore, 865
lesser cornstalk borer, 865
pest management, 865
biological control, 51, 81, 282, 298, 317, 448, 508, 657, 669, 697, 896, 925, 989, 1184, 1364, 1417, 1454, 1527, 1653
Agrilus planipennis, 925
Aonidiella aurantii, 51
Aphis glycines, 282, 1417
apple, 1364
Asopinae, 1454
augmentation, 1527
Beauveria bassiana, 298
Bemisia tabaci, 669
Binodoxys communis, 282, 1417
Braconidae, 657
Brassicaceae, 1454
Carmenta ithacae, 317
climate change, 1653
climate match, 448
CLIMEX model, 317
competition, 896
cover crop, 298
Curculio caryae, 298
Delphastus catalinae, 669
development, 989
Diadegma insulare, 1653
diet, 669
ectoparasitoid, 697
ELISA, 1527
Eriosoma lanigerum, 1364
fluctuating temperature, 1653
foraging behavior, 508
habitat structure, 508
Habrobracon hebetor, 697
host-plant resistance, 282, 1417
host plant use, 81
interspecific competition, 925
invasive species, 81
Laricobius, 896
Lepidoptera, 81
longevity, 51
mealybug, 1184
natural enemies, 1364
Noctuidae, 657
parasitism, 1184
parasitoids, 1184
Parthenium hysterophorus, 317
pecan, 298
phenology, 448
photoperiod, 697
Plutella xylostella, 1653
predation, 896
predation monitoring, 1527
predator, 669
prey density, 508
Prokelisia marginata, 448
protein marking, 1527
reproduction, 989
reproductive diapause, 697
Sesiidae, 317
sex ratio, 51
size, 51
southern Africa, 989
soybean, 282, 1417
Spartina alterniflora, 448
Spathius galinae, 925
Spodoptera exigua, 657
survey, 1364
temperature, 989
Tetrastichus planipennisi, 925
Totricidae, 81
tritrophic interactions, 508
biological invasion,
host plant, 1398
Indian Oceanic Islands, 1398
Kerguelen Islands, 1398
Myzus ascalonicus, 1398
biology,
fertility life table, 849
Lepidoptera, 849
oriental fruit moth, 849
biotic interactions,
Aphaenogaster picea, 463
Aphaenogaster rudis, 463
community ecology, 463
Lasius alienus, 463
Black Hills,
Carabidae, 1069
Coleoptera, 1069
ground beetles, 1069
vegetation, 1069
Blissus insularis Barber,
aggregation, 887
host-plant volatiles, 887
pheromones, 887
Y-tube olfactometer, 887
blue-stain,
Dendroctonus, 478
nitrogen, 478
pine, 478
symbiosis, 478
blueberry gall midge,
monitoring, 455
phenology, 455
trap, 455
blueberry maggot,
compost, 370
fly emergence, 370
pine needles, 370
Rhagoletis mendax, 370
body mass,
development rate, 714
fitness, 714
intrinsic rate of increase, 714
SSI model, 714
body size,
diapause, 1612
hibernation, 1612

natural selection, 1612
polyandry, 1612
Bombus, 905, 1332
clover, 905
competition, 1332
corolla length, 1332
elevation, 1332
microsatellites, 905
pollinators, 905
proboscis length, 1332
bottleneck,
F_{ST}, 188
genetic fingerprinting, 188
genetic structure, 188
Tribolium castaneum, 188
bottom-up effects,
detritus, 1062
longevity, 1062
mosquitoes, 1062
Braconidae, 587, 657
activity, 587
biological control, 657
Ichneumonidae, 587
Noctuidae, 657
relative humidity, 587
Spodoptera exigua, 657
temperature, 587
Brassicaceae,
Asopinae, 1454
biological control, 1454
Brevicoryne brassicae,
bacterial community, 1386
endosymbiont, 1386
real-time qPCR, 1386
16S ribosomal DNA, 1386
Brontispa longissima,
fecundity, 1208
growth and development, 1208
host plant, 1208
life table, 1208
Bt corn, 200, 1268
arthropod abundance, 1268
monitoring, 1268
PRC analysis, 1268
resistance management, 200
riparian area, 1268
simulation, 200
Bt cowpea,
resistance management, 1255
simulation, 1255
bumble bee,
floral volatile, 855
greenhouse, 855
repellent, 855
tomato, 855
Burg El-Arab,
Egypt, 11
Metarhizium anisopliae, 11
monthly abundance, 11
sand flies, 11

C

cannibalism, 166, 1501
disease, 166
flour conditioning, 1501
katydid, 166
lysozyme, 166
melanism, 166
population dynamics, 1501
Tribolium castaneum, 1501
canola,
epigeic predators, 72
intercropping, 72
oilseed rape, 72
sustainable agriculture, 72
Capsicum anuum,
conservation biological control, 612
European corn borer, 612
intercropping, 612
Carabidae, 1069, 1097
Black Hills, 1069
Coleoptera, 1069
compost, 1097
ground beetles, 1069
highbush blueberries, 1097
mulch, 1097
Staphylinidae, 1097
vegetation, 1069
Cardinium,
cytoplasmic incompatibility, 833
infection frequency, 833
Sogatella furcifera, 833
Wolbachia, 833
Carmenta ithacae,
biological control, 317
CLIMEX model, 317
Parthenium hysterophorus, 317
Sesiidae, 317
Carposina sasakii,
duration, 349
fecundity, 349
host plant, 349
survival rate, 349
cattle,
Bailey's triple catch, 20
fluorescent marking, 20
forecasting, 20
longevity, 20
cDNA,
ecotype, 1008
genomic DNA, 1008
Nylanderia pubens, 1008
Partrechina pubens, 1008
Cerambycidae, 1350, 1587
host finding, 1350
host volatile, 1587
Ips, 1350
Monochamus, 1587
pheromone, 1587
saproxylic, 1350
Scolytinae, 1350
2-(undecyloxy)-ethanol, 1587
Cerotoma trifurcata,
Aphis glycines, 750
Empoasca fabae, 750
soybean, 750
winter rye, 750
Ceutorhynchus alliariae,
Ceutorhynchus roberti, 1086
competitive coexistence, 1086
niche overlap, 1086
niche partitioning, 1086
Ceutorhynchus roberti,
Ceutorhynchus alliariae, 1086
competitive coexistence, 1086
niche overlap, 1086
niche partitioning, 1086
chemical defense,
deciduous forest, 541
herbivores, 541
insects, 541
ontogenetic succession, 541
chemical ecology,
fungal endophytes, 433
gall induction, 433
infection, 433
insects, 433
chestnut volatile organic compound profile,
Curculio, 933
electroantennogram, 933
gas chromatography-mass spectrometry, 933
solid phase microextraction, 933
Choristoneura rosaceana,
exposure, 1481
Grapholita molesta, 1481
pheromone, 1481
sexual behavior, 1481
Chrysodeix includes,
Glycine max, 551
induced resistance, 551
jasmonic acid, 551
Circulifer,
beet leafhopper, 768
nonparametric multiplicative regression, 768
pest management, 768
purple top disease, 768
classical biological control,
Ailanthus altissima, 118
Eucryptorrhynchus brandti, 118
host-specificity, 118
invasive species, 118
climate change,
biological control, 1653
Diadegma insulare, 1653
fluctuating temperature, 1653
Plutella xylostella, 1653
climate match,
biological control, 448
phenology, 448
Prokelisia marginata, 448
Spartina alterniflora, 448

CLIMEX model,
biological control, 317
Carmenta ithacae, 317
Parthenium hysterophorus, 317
Sesiidae, 317
clover,
Bombus, 905
microsatellites, 905
pollinators, 905
coccid,
alkaloids, 333
coffee plants, 333
nutrients, 333
phenolic compounds, 333
Coccinella septempunctata,
fat body, 676
Harmonia axyridis, 676
temperature, 676
weight, 676
coffee,
agroforestry, 1107
Azteca instabilis, 1107
Pseudacteon, 1107
trait-meditated interactions, 1107
coffee plants,
alkaloids, 333
coccid, 333
nutrients, 333
phenolic compounds, 333
CO-I,
cryptic species, 1239
DNA barcoding, 1239
mirids, 1239
species-specific marker, 1239
Coleoptera, 392, 1069
Black Hills, 1069
Carabidae, 1069
Curculionidae, 392
ground beetles, 1069
Scolytinae, 392
secondary metabolites, 392
terpenes, 392
vegetation, 1069
Colorado potato beetle,
crop rotation, 1553
diapause, 1553
geographic information system, 1553
integrated pest management, 1553
common chickweed,
glyphosate, 362
paraquat, 362
Stellaria media, 362
tobacco thrips, 362
community ecology,
Aphaenogaster picea, 463
Aphaenogaster rudis, 463
biotic interactions, 463
Lasius alienus, 463
competition, 896, 1332
biological control, 896
Bombus, 1332
corolla length, 1332
elevation, 1332
Laricobius, 896
predation, 896
proboscis length, 1332
competitive coexistence,
Ceutorhynchus alliariae, 1086
Ceutorhynchus roberti, 1086
niche overlap, 1086
niche partitioning, 1086
compost, 370, 1097
blueberry maggot, 370
Carabidae, 1097
fly emergence, 370
highbush blueberries, 1097
mulch, 1097
pine needles, 370
Rhagoletis mendax, 370
Staphylinidae, 1097
conifers,
ancestral state reconstruction, 723
bark beetles, 723
Dendroctonus, 723
outbreaks, 723
conservation biological control, 275, 612, 621
Capsicum anuum, 612
energy sugarcane, 275
European corn borer, 612
Humulus lupulus, 621
intercropping, 612
pest resurgence, 621
Poaceae, 275
Podosphaera macularis, 621
refuge, 275
stemborer, 275
conspecifics,
host fruit, 387
2-methyl-6-vinylpyrazine, 387
papaya fruit fly, 387
pheromone, 387
copulatory behavior,
courtship, 562
daily rhythm, 562
pheromones, 562
semiochemicals, 562
corn earworm,
lower lethal temperature, 172
lower lethal time, 172
supercooling point, 172
corolla length,
Bombus, 1332
competition, 1332
elevation, 1332
proboscis length, 1332
Costa Rica,
entomological net, 1544
immature stages, 1544
Malaise trap, 1544
phytotelmata, 1544
cotton, 125, 342
Gossypium, 342
Hemiptera, 125
horsemint, 125
Lygus, 342
olfactometer, 125
prior experience, 125
western tarnished plant bug, 342
within-plant distribution, 342
courtship,
copulatory behavior, 562
daily rhythm, 562
pheromones, 562
semiochemicals, 562
cover crop,
Beauveria bassiana, 298
biological control, 298
Curculio caryae, 298
pecan, 298
cricket,
feeding, 1644
Gryllus, 1644
orthopteran, 1644
reproduction, 1644
crop rotation,
Colorado potato beetle, 1553
diapause, 1553
geographic information system, 1553
integrated pest management, 1553
Cry3Bb1,
Bacillus thuringiensis, 1037
nontarget risk assessment, 1037
Theridion impressum, 1037
cryptic species,
CO-I, 1239
DNA barcoding, 1239
mirids, 1239
species-specific marker, 1239
cucumber,
pan trapping, 532
wild bees, 532
Wisconsin, 532
cultural control,
Bactrocera oleae, 1177
fruit fly dispersal, 1177
mature larvae, 1177
teneral adult, 1177
Curculio,
chestnut volatile organic compound profile, 933
electroantennogram, 933
gas chromatography-mass spectrometry, 933
solid phase microextraction, 933

Curculio caryae,
Beauveria bassiana, 298
biological control, 298
cover crop, 298
pecan, 298
Curculionidae,
Coleoptera, 392
Scolytinae, 392
secondary metabolites, 392
terpenes, 392
Cybocephalus spp.,
Encarsia citrina, 377
euonymus scale (*Unaspis euonymi*), 377
nontarget effects, 377
nursery production, 377
Cydia pomonella, 399, 407
apple, 399, 407
mating disruption, 399, 407
monitoring, 399
plant volatiles, 407
sex pheromone, 399
traps, 407
cytoplasmic incompatibility,
Cardinium, 833
infection frequency, 833
Sogatella furcifera, 833
Wolbachia, 833

D

daily rhythm,
copulatory behavior, 562
courtship, 562
pheromones, 562
semiochemicals, 562
day collections,
Araneae, 1534
Atlantic Rainforest, 1534
night collections, 1534
sampling effort, 1534
deciduous forest,
chemical defense, 541
herbivores, 541
insects, 541
ontogenetic succession, 541
degree-day, 761, 950
developmental rate, 950
integrated pest management, 761
nonbiting midge, 950
Oulema melanopus, 761
predictive model, 761
sampling, 761
thermal threshold, 950
toxicity test organism, 950
Deladenus siricidicola,
Ibalia leucospoides, 289
natural enemies, 289
parasitic nematode, 289
parasitoids, 289
Delia radicum,
root-feeding insect, 159
soil moisture, 159
soil temperature, 159
Delphacidae,
global warming, 1231
mate-location behavior, 1231
planthopper, 1231
temperature, 1231
Delphastus catalinae,
Bemisia tabaci, 669
biological control, 669
diet, 669
predator, 669
Dendroctonus, 478, 723
ancestral state reconstruction, 723
bark beetles, 723
blue-stain, 478
conifers, 723
nitrogen, 478
outbreaks, 723
pine, 478
symbiosis, 478
detritus,
bottom-up effects, 1062
longevity, 1062
mosquitoes, 1062
development, 1, 706, 989, 1190
age-specific life table, 1
Bactericera cockerelli, 1190
biological control, 989
Echinothrips americanus, 706
fecundity, 1
Lygus lineolaris, 1
parthenogenesis, 706
reproduction, 989
sexual reproduction, 706
southern Africa, 989
temperature, 1, 989, 1190
tomato psyllid (potato psyllid), 1190
development rate,
body mass, 714
fitness, 714
intrinsic rate of increase, 714
SSI model, 714
development time,
diapause, 415
plant bug, 415
rearing, 415
developmental rate,
degree-day, 950
nonbiting midge, 950
thermal threshold, 950
toxicity test organism, 950
Diadegma insulare,
biological control, 1653
climate change, 1653
fluctuating temperature, 1653
Plutella xylostella, 1653
diapause, 415, 1199, 1302, 1553, 1606, 1612, 1621
Adelphocoris suturalis, 1606
Apolygus lucorum, 1606
body size, 1612
Colorado potato beetle, 1553
crop rotation, 1553
development time, 415
flight, 1199
geographic information system, 1553
hibernation, 1612
integrated pest management, 1553
Loxostege sticticalis, 1199
Lygus lineolaris, 1302
natural selection, 1612
Osmia rufa, 1621
photoperiod, 1606
plant bug, 415
plant debris, 1302
polyandry, 1612
rearing, 415
reproduction, 1199
sensitive stage, 1606
simulated climate change, 1621
tarnished plant bug, 1302
temperature, 1621
winter host plants, 1302
Diatraea saccharalis,
host-plant resistance, 571
Oryza sativa, 571
oviposition behavior, 571
diel activity,
diurnal, 594
intracrown, 594
Pikonema alaskensis, 594
thermoregulation, 594
diet,
Bemisia tabaci, 669
biological control, 669
Delphastus catalinae, 669
predator, 669
disease,
cannibalism, 166
katydid, 166
lysozyme, 166
melanism, 166
dispersal, 578, 1050, 1466
biocontrol, 1466
geostatistics, 1050
Gomphocerinae, 1050
harmonic radar, 578
Hylobius warreni, 578
Italian locust, 1050
movement, 578
orientation, 1466
Persicaria perfoliata, 1466
Rhinoncomimus latipes, 1466
sampling, 578
spatial pattern, 1050

distribution,
Apis mellifera, 1462
phenology, 1462
swarm, 1462
Diuraphis noxia,
agriculture, 1505
pest, 1505
Russian wheat aphid, 1505
spatial model, 1505
diurnal,
diel activity, 594
intracrown, 594
Pikonema alaskensis, 594
thermoregulation, 594
diversity,
agricultural areas, 776
flight pattern, 776
Xyleborini ambrosia beetles, 776
DNA barcoding,
CO-I, 1239
cryptic species, 1239
mirids, 1239
species-specific marker, 1239
dog-strangling vine,
fecundity, 841
host range, 841
Hypena, 841
larval development, 841
Doryctobracon,
Aganaspis, 233
Anastrepha, 233
Asobara, 233
Neosilba, 233
dose–response,
emerald ash borer, 648
lactone, 648
pheromone, 648
(3Z)-dodecen-12-olide, 648
drainage ditch,
biodiversity, 802
macroinvertebrate, 802
nutrients, 802
water quality, 802
dung beetle assemblages,
abiotic factors, 785
Mediterranean bioclimate, 785
spatial scale, 785
temporal scale, 785
duration,
Carposina sasakii, 349
fecundity, 349
host plant, 349
survival rate, 349

E

Echinothrips americanus,
development, 706
parthenogenesis, 706
sexual reproduction, 706
ecosystem services,
mid-Atlantic agriculture, 813
native bees, 813
non-*Apis* bee crop pollination, 813
wild bees, 813
ecotype,
cDNA, 1008
genomic DNA, 1008
Nylanderia pubens, 1008
Partrechina pubens, 1008
ectoparasitoid,
biological control, 697
Habrobracon hebetor, 697
photoperiod, 697
reproductive diapause, 697
egg clustering,
adult emergence, 971
egg desiccation hypothesis, 971
egg eclosion, 971
waste management, 971
egg desiccation hypothesis,
adult emergence, 971
egg clustering, 971
egg eclosion, 971
waste management, 971
egg diapause,
hatch, 1662
Scymnus camptodromus, 1662
temperature, 1662
egg eclosion,
adult emergence, 971
egg clustering, 971
egg desiccation hypothesis, 971
waste management, 971
egg masses,
hydrological seasonality, 1322
macroinvertebrates, 1322
recruitment, 1322
tropical streams, 1322
Egypt,
Burg El-Arab, 11
Metarhizium anisopliae, 11
monthly abundance, 11
sand flies, 11
Eisenia fetida,
bermudagrass, 1139
feeding ecology, 1139
plant insect interactions, 1139
Scapteriscus, 1139
elaiosome,
ant–plant interactions, 1405
mutualism, 1405
myrmecochory, 1405
trophic interactions, 1405
electrical penetration graph,
Homalodisca vitripennis, 1215
ingestion, 1215
insect feeding, 1215
temperature, 1215
electroantennogram,
chestnut volatile organic compound profile, 933
Curculio, 933
gas chromatography-mass spectrometry, 933
solid phase microextraction, 933
electroantennography,
manuka oil, 1597
phoebe oil, 1597
redbay ambrosia beetle, 1597
silkbay, 1597
elevation,
Bombus, 1332
competition, 1332
corolla length, 1332
proboscis length, 1332
ELISA,
augmentation, 1527
biological control, 1527
predation monitoring, 1527
protein marking, 1527
emerald ash borer,
dose–response, 648
lactone, 648
pheromone, 648
(3Z)-dodecen-12-olide, 648
Empoasca fabae,
Aphis glycines, 750
Cerotoma trifurcata, 750
soybean, 750
winter rye, 750
encapsulation,
apoptosis, 688
Galleria mellonella, 688
gibberellic acid, 688
hemocyte, 688
Encarsia citrina,
Cybocephalus spp., 377
euonymus scale (*Unaspis euonymi*), 377
nontarget effects, 377
nursery production, 377
endosymbiont,
bacterial community, 1386
Brevicoryne brassicae, 1386
real-time qPCR, 1386
16S ribosomal DNA, 1386
energy sugarcane,
conservation biological control, 275
Poaceae, 275
refuge, 275
stemborer, 275
entomological net,
Costa Rica, 1544
immature stages, 1544
Malaise trap, 1544
phytotelmata, 1544

entomopathogen,
Entomophaga maimaiga, 1115
fungal epizootic, 1115
gypsy moth, 1115
soil pH, 1115
entomopathogenic fungi,
immunity, 959
nitric oxide, 959
phenol oxidases, 959
scarabaeidae larvae, 959
Entomophaga maimaiga,
entomopathogen, 1115
fungal epizootic, 1115
gypsy moth, 1115
soil pH, 1115
environmental condition,
adult trees, 828
pine wood nematode, 828
shading, 828
virulence, 828
environmental risk assessment,
genetically engineered plants, 1687
ladybird beetles, 1687
nontarget arthropods, 1687
environmental stress,
geometric morphometrics, 979
microhabitat, 979
preadaptation, 979
epigeic predators,
canola, 72
intercropping, 72
oilseed rape, 72
sustainable agriculture, 72
Eriosoma lanigerum,
apple, 1364
biological control, 1364
natural enemies, 1364
survey, 1364
ethanol, 636, 865
ambrosia beetles, 636
bioenergy, 865
herbivore, 865
lesser cornstalk borer, 865
pest management, 865
solid phase microextraction, 636
Xylosandrus germanus, 636
Eucryptorrhynchus brandti,
Ailanthus altissima, 118
classical biological control, 118
host-specificity, 118
invasive species, 118
euonymus scale (*Unaspis euonymi*),
Cybocephalus spp., 377
Encarsia citrina, 377
nontarget effects, 377
nursery production, 377
European corn borer,
Capsicum anuum, 612
conservation biological control, 612
intercropping, 612
exclusion experiments,
female reproductive success, 1163
foraging behavior, 1163
pollen loads, 1163
exotic,
bean plataspid, 40
host preference, 40
invasive, 40
kudzu bug, 40
exotic species,
feeding preference, 238
niche partitioning, 238
Onthophagus, 238
Phanaeus, 238
exotics and invasive species,
Agrilus planipennis, 245
indigenous, 245
parasitoids, 245
wood borers, 245
exposure,
Choristoneura rosaceana, 1481
Grapholita molesta, 1481
pheromone, 1481
sexual behavior, 1481

F

fall armyworm,
maize, 1131
Spodoptera frugiperda, 1131
sweet corn, 1131
false rings,
Adelges tsugae, 523
Fiorinia externa, 523
herbivory, 523
Tsuga canadensis, 523
fat body,
Coccinella septempunctata, 676
Harmonia axyridis, 676
temperature, 676
weight, 676
F_{ST},
bottleneck, 188
genetic fingerprinting, 188
genetic structure, 188
Tribolium castaneum, 188
fecundity, 1, 349, 841, 1208
age-specific life table, 1
Brontispa longissima, 1208
Carposina sasakii, 349
development, 1
dog-strangling vine, 841
duration, 349
growth and development, 1208
host plant, 349, 1208
host range, 841
Hypena, 841
larval development, 841
life table, 1208
Lygus lineolaris, 1
survival rate, 349
temperature, 1
feeding,
cricket, 1644
Gryllus, 1644
orthopteran, 1644
reproduction, 1644
feeding ecology,
bermudagrass, 1139
Eisenia fetida, 1139
plant insect interactions, 1139
Scapteriscus, 1139
feeding preference,
exotic species, 238
niche partitioning, 238
Onthophagus, 238
Phanaeus, 238
female reproductive success,
exclusion experiments, 1163
foraging behavior, 1163
pollen loads, 1163
fertility life table,
biology, 849
Lepidoptera, 849
oriental fruit moth, 849
Fiorinia externa,
Adelges tsugae, 523
false rings, 523
herbivory, 523
Tsuga canadensis, 523
fitness,
body mass, 714
development rate, 714
intrinsic rate of increase, 714
SSI model, 714
flight,
diapause, 1199
Loxostege sticticalis, 1199
reproduction, 1199
flight pattern,
agricultural areas, 776
diversity, 776
Xyleborini ambrosia beetles, 776
floral volatile,
bumble bee, 855
greenhouse, 855
repellent, 855
tomato, 855
flour conditioning,
cannibalism, 1501
population dynamics, 1501
Tribolium castaneum, 1501
flower diversity,
aphids, 1077
natural enemy diversity, 1077
natural pest suppression, 1077

fluctuating temperature,
biological control, 1653
climate change, 1653
Diadegma insulare, 1653
Plutella xylostella, 1653
fluorescent marking,
Bailey's triple catch, 20
cattle, 20
forecasting, 20
longevity, 20
fly emergence,
blueberry maggot, 370
compost, 370
pine needles, 370
Rhagoletis mendax, 370
foraging behavior, 508, 516, 1163
biological control, 508
exclusion experiments, 1163
female reproductive success, 1163
habitat structure, 508
movement, 516
partial refugia, 516
pollen loads, 1163
prey density, 508
tritrophic interactions, 508, 516
forecasting,
Bailey's triple catch, 20
cattle, 20
fluorescent marking, 20
longevity, 20
forest decline,
Betula, 1372
host plant resistance, 1372
insect outbreaks, 1372
population dynamics, 1372
Formicidae,
ant diversity, 222
lowbush blueberry, 222
Maine, 222
foxglove aphid,
Aulacorthum solani, 665
host suitability, 665
swallow-wort, 665
Vincetoxicum, 665
Frankliniella,
banker plants, 487
minute pirate bugs, 487
Orius, 487
thrips, 487
fruit fly dispersal,
Bactrocera oleae, 1177
cultural control, 1177
mature larvae, 1177
teneral adult, 1177
fungal endophytes,
chemical ecology, 433
gall induction, 433
infection, 433
insects, 433
fungal epizootic,
entomopathogen, 1115
Entomophaga maimaiga, 1115
gypsy moth, 1115
soil pH, 1115
funnel traps,
Pinus, 91
Sirex, 91
Tremex, 91
Urocerus, 91
Fusarium solani,
beetle–fungi association, 255
phoretic, 255
Scolytogenes birosimensis, 255
tree mortality, 255
Fusarium,
intraspecies mixture, 1029
pest, 1029
predator, 1029
refuge, 1029

G

gall induction,
chemical ecology, 433
fungal endophytes, 433
infection, 433
insects, 433
Galleria mellonella,
apoptosis, 688
encapsulation, 688
gibberellic acid, 688
hemocyte, 688
gas chromatography-mass spectrometry,
chestnut volatile organic compound profile, 933
Curculio, 933
electroantennogram, 933
solid phase microextraction, 933
genetic fingerprinting,
bottleneck, 188
F_{ST}, 188
genetic structure, 188
Tribolium castaneum, 188
genetic structure,
bottleneck, 188
F_{ST}, 188
genetic fingerprinting, 188
Tribolium castaneum, 188
genetically engineered plants,
environmental risk assessment, 1687
ladybird beetles, 1687
nontarget arthropods, 1687
genomic DNA,
cDNA, 1008
ecotype, 1008
Nylanderia pubens, 1008
Partrechina pubens, 1008
geographic information system,
Colorado potato beetle, 1553
crop rotation, 1553
diapause, 1553
integrated pest management, 1553
geometric morphometrics,
environmental stress, 979
microhabitat, 979
preadaptation, 979
geostatistics,
dispersal, 1050
Gomphocerinae, 1050
Italian locust, 1050
spatial pattern, 1050
gibberellic acid,
apoptosis, 688
encapsulation, 688
Galleria mellonella, 688
hemocyte, 688
glassy-winged sharpshooter,
insect dispersal, 1279
mark–capture, 1279
Pierce's disease, 1279
Xylella fastidiosa, 1279
global change,
jasmonic acid, 1435
plant–insect interactions, 1435
salicylic acid, 1435
global warming,
Delphacidae, 1231
mate-location behavior, 1231
planthopper, 1231
temperature, 1231
Glycine max,
Chrysodeix includes, 551
induced resistance, 551
jasmonic acid, 551
glyphosate,
common chickweed, 362
paraquat, 362
Stellaria media, 362
tobacco thrips, 362
Gomphocerinae,
dispersal, 1050
geostatistics, 1050
Italian locust, 1050
spatial pattern, 1050
Gossypium,
cotton, 342
Lygus, 342
western tarnished plant bug, 342
within-plant distribution, 342
Grapholita molesta,
Choristoneura rosaceana, 1481
exposure, 1481
pheromone, 1481
sexual behavior, 1481
green leaf volatiles,
Pinus albicaulis, 1575
Pinus contorta, 1575
semiochemicals, 1575
verbenone, 1575

greenhouse,
bumble bee, 855
floral volatile, 855
repellent, 855
tomato, 855
ground beetles,
Black Hills, 1069
Carabidae, 1069
Coleoptera, 1069
vegetation, 1069
growth and development,
Brontispa longissima, 1208
fecundity, 1208
host plant, 1208
life table, 1208
Gryllus,
cricket, 1644
feeding, 1644
orthopteran, 1644
reproduction, 1644
gypsy moth,
entomopathogen, 1115
Entomophaga maimaiga, 1115
fungal epizootic, 1115
soil pH, 1115

H

habitat,
autecology, 966
integrated pest management (IPM), 966
soil-dwelling pest, 966
habitat structure,
biological control, 508
foraging behavior, 508
prey density, 508
tritrophic interactions, 508
Habrobracon hebetor,
biological control, 697
ectoparasitoid, 697
photoperiod, 697
reproductive diapause, 697
Harmonia axyridis, 676, 1169, 1565
Coccinella septempunctata, 676
fat body, 676
integrated pest management, 1565
olfactometer, 1169
Orius insidiosus, 1565
potassium metabisulfite, 1169
repellent, 1169
soybean aphid, 1565
spirotetramat, 1565
temperature, 676
vineyard, 1169
weight, 676
harmonic radar,
dispersal, 578
Hylobius warreni, 578
movement, 578
sampling, 578
hatch,
egg diapause, 1662
Scymnus camptodromus, 1662
temperature, 1662
Helicoverpa armigera,
artificial diet, 30
life table, 30
Zea mays, 30
Hemiptera,
cotton, 125
horsemint, 125
olfactometer, 125
prior experience, 125
hemocyte,
apoptosis, 688
encapsulation, 688
Galleria mellonella, 688
gibberellic acid, 688
herbivore, 541, 865, 872
apple orchard, 872
aromatic plant, 872
arthropod community, 872
bioenergy, 865
chemical defense, 541
deciduous forest, 541
ethanol, 865
insects, 541
lesser cornstalk borer, 865
ontogenetic succession, 541
pest management, 865
predator, 872
herbivory,
Adelges tsugae, 523
false rings, 523
Fiorinia externa, 523
Tsuga canadensis, 523
hibernation,
body size, 1612
diapause, 1612
natural selection, 1612
polyandry, 1612
high-temperature stress,
Bemisia tabaci biotype B, 1672
life table, 1672
long-term fitness effects, 1672
highbush blueberries,
Carabidae, 1097
compost, 1097
mulch, 1097
Staphylinidae, 1097
Homalodisca vitripennis,
electrical penetration graph, 1215
ingestion, 1215
insect feeding, 1215
temperature, 1215
Homoptera,
aphid, 59
invasive ant, 59
myrmecophily, 59
Myrmica rubra, 59
honey bees,
Aethina tumida, 152
insect trapping, 152
small hive beetle larvae, 152
horsemint,
cotton, 125
Hemiptera, 125
olfactometer, 125
prior experience, 125
host finding,
Cerambycidae, 1350
Ips, 1350
saproxylic, 1350
Scolytinae, 1350
host fruit,
conspecifics, 387
papaya fruit fly, 387
pheromone, 387
2-methyl-6-vinylpyrazine, 387
host plant, 349, 1208, 1398
biological invasion, 1398
Brontispa longissima, 1208
Carposina sasakii, 349
duration, 349
fecundity, 349, 1208
growth and development, 1208
Indian Oceanic Islands, 1398
Kerguelen Islands, 1398
life table, 1208
Myzus ascalonicus, 1398
survival rate, 349
host plant resistance,
Betula, 1372
forest decline, 1372
insect outbreaks, 1372
population dynamics, 1372
host plant use,
biological control, 81
invasive species, 81
Lepidoptera, 81
Totricidae, 81
host preference,
bean plataspid, 40
exotic, 40
invasive, 40
kudzu bug, 40
host range,
dog-strangling vine, 841
fecundity, 841
Hypena, 841
larval development, 841
host suitability, 665, 1125
Aulacorthum solani, 665
Bemisia tabaci B biotype, 1125
foxglove aphid, 665
interspecific interaction, 1125
leaf age, 1125
swallow-wort, 665
Trialeurodes vaporariorum, 1125
Vincetoxicum, 665

host volatile,
Cerambycidae, 1587
Monochamus, 1587
pheromone, 1587
2-(undecyloxy)-ethanol, 1587
host-associated divergence,
Asphondylia borrichiae, 1246
host-race formation, 1246
sympatric speciation, 1246
host-plant resistance, 282, 571, 1417, 1426
Aphis glycines, 282, 1417, 1426
Binodoxys communis, 282, 1417
biological control, 282, 1417
Diatraea saccharalis, 571
movement, 1426
Oryza sativa, 571
oviposition behavior, 571
soybean, 282, 1417
within-plant distribution, 1426
host-plant volatiles,
aggregation, 887
Blissus insularis Barber, 887
pheromones, 887
Y-tube olfactometer, 887
host-race formation,
Asphondylia borrichiae, 1246
host-associated divergence, 1246
sympatric speciation, 1246
host-specificity,
Ailanthus altissima, 118
classical biological control, 118
Eucryptorrhynchus brandti, 118
invasive species, 118
Humulus lupulus,
conservation biological control, 621
pest resurgence, 621
Podosphaera macularis, 621
hydrological seasonality,
egg masses, 1322
macroinvertebrates, 1322
recruitment, 1322
tropical streams, 1322
Hylobius warreni,
dispersal, 578
harmonic radar, 578
movement, 578
sampling, 578
Hypena,
dog-strangling vine, 841
fecundity, 841
host range, 841
larval development, 841

I

Ibalia leucospoides,
Deladenus siricidicola, 289
natural enemies, 289
parasitic nematode, 289
parasitoids, 289
Ibalia,
Megarhyssa, 731
Pseudorhyssa, 731
Rhyssa, 731
Sirex noctilio, 731
Ichneumonidae,
activity, 587
Braconidae, 587
relative humidity, 587
temperature, 587
immature stages,
Costa Rica, 1544
entomological net, 1544
Malaise trap, 1544
phytotelmata, 1544
immunity,
entomopathogenic fungi, 959
nitric oxide, 959
phenol oxidases, 959
scarabaeidae larvae, 959
Indian Oceanic Islands,
biological invasion, 1398
host plant, 1398
Kerguelen Islands, 1398
Myzus ascalonicus, 1398
indigenous,
Agrilus planipennis, 245
exotics and invasive species, 245
parasitoids, 245
wood borers, 245
indirect interactions,
induced defenses, 1409
leaf traits, 1409
terpenes, 1409
vertebrate herbivory, 1409
induced defenses,
indirect interactions, 1409
leaf traits, 1409
terpenes, 1409
vertebrate herbivory, 1409
induced resistance,
Chrysodeix includes, 551
Glycine max, 551
jasmonic acid, 551
infection,
chemical ecology, 433
fungal endophytes, 433
gall induction, 433
insects, 433
infection frequency,
Cardinium, 833
cytoplasmic incompatibility, 833
Sogatella furcifera, 833
Wolbachia, 833
ingestion,
electrical penetration graph, 1215
Homalodisca vitripennis, 1215
insect feeding, 1215
temperature, 1215
insect dispersal,
glassy-winged sharpshooter, 1279
mark–capture, 1279
Pierce's disease, 1279
Xylella fastidiosa, 1279
insect feeding,
electrical penetration graph, 1215
Homalodisca vitripennis, 1215
ingestion, 1215
temperature, 1215
insect outbreaks,
Betula, 1372
forest decline, 1372
host plant resistance, 1372
population dynamics, 1372
insect trapping,
Aethina tumida, 152
honey bees, 152
small hive beetle larvae, 152
insecticide residue,
oviposition preference, 880
Plutella xylostella, 880
resistance management, 880
transgenic plants, 880
insects, 433, 541
chemical defense, 541
chemical ecology, 433
deciduous forest, 541
fungal endophytes, 433
gall induction, 433
herbivores, 541
infection, 433
ontogenetic succession, 541
integrated pest management, 761, 966, 1553, 1565
autecology, 966
Colorado potato beetle, 1553
crop rotation, 1553
degree-days, 761
diapause, 1553
geographic information system, 1553
habitat, 966
Harmonia axyridis, 1565
Orius insidiosus, 1565
Oulema melanopus, 761
predictive model, 761
sampling, 761
soil-dwelling pest, 966
soybean aphid, 1565
spirotetramat, 1565
intercropping, 72, 612
canola, 72
Capsicum anuum, 612
conservation biological control, 612
epigeic predators, 72
European corn borer, 612
oilseed rape, 72
sustainable agriculture, 72

interspecific competition,
Agrilus planipennis, 925
biological control, 925
Spathius galinae, 925
Tetrastichus planipennisi, 925
interspecific interaction,
Bemisia tabaci B biotype, 1125
host suitability, 1125
leaf age, 1125
Trialeurodes vaporariorum, 1125
intracrown,
diel activity, 594
diurnal, 594
Pikonema alaskensis, 594
thermoregulation, 594
intraspecies mixture,
Fusarium, 1029
pest, 1029
predator, 1029
refuge, 1029
intrinsic rate of increase,
body mass, 714
development rate, 714
fitness, 714
SSI model, 714
invasion ecology,
artificial diet, 265
olive fruit fly, 265
sterile insect technology, 265
symbiosis, 265
invasive,
bean plataspid, 40
exotic, 40
host preference, 40
kudzu bug, 40
invasive ant,
aphid, 59
Homoptera, 59
myrmecophily, 59
Myrmica rubra, 59
invasive pest,
ambrosia beetle, 822
artificial diet, 822
laboratory rearing, 822
invasive species, 81, 118, 1680
Ailanthus altissima, 118
biological control, 81
classical biological control, 118
Eucryptorrhynchus brandti, 118
host plant use, 81
host-specificity, 118
Isoptera, 1680
Lepidoptera, 81
population genetics, 1680
Reticulitermes, 1680
social insect, 1680
Totricidae, 81
Ips,
Cerambycidae, 1350
host finding, 1350
saproxylic, 1350
Scolytinae, 1350
Iran,
beet armyworm, 326
life table, 326
reproduction, 326
soybean, 326
irradiated host,
Bactrocera oleae (Rossi), 497
larval parasitoid, 497
longevity, 497
parasitism, 497
Isoptera,
invasive species, 1680
population genetics, 1680
Reticulitermes, 1680
social insect, 1680
Italian locust,
dispersal, 1050
geostatistics, 1050
Gomphocerinae, 1050
spatial pattern, 1050

J

jasmonic acid, 551, 1435, 1443
Bemisia tabaci, 1443
Chrysodeix includes, 551
global change, 1435
Glycine max, 551
induced resistance, 551
plant–insect interactions, 1435
salicylic acid, 1435, 1443
tomato, 1443
whitefly preconditioning, 1443
Jolly–Seber's stochastic method,
Bailey's triple-catch, 141
loss rate, 141
Monte-Carlo simulation, 141
population estimates, 141

K

kairomone,
arrestment, 603
larvae, 603
microencapsulated, 603
pear ester, 603
katydid,
cannibalism, 166
disease, 166
lysozyme, 166
melanism, 166
Kerguelen Islands,
biological invasion, 1398
host plant, 1398
Indian Oceanic Islands, 1398
Myzus ascalonicus, 1398
kudzu bug,
bean plataspid, 40
exotic, 40
host preference, 40
invasive, 40

L

laboratory rearing,
ambrosia beetle, 822
artificial diet, 822
invasive pest, 822
lactone,
dose–response, 648
emerald ash borer, 648
pheromone, 648
(3Z)-dodecen-12-olide, 648
ladybird beetles,
environmental risk assessment, 1687
genetically engineered plants, 1687
nontarget arthropods, 1687
Lambdina fiscellaria,
seasonal ecology, 1290
thermal requirement, 1290
thermal response, 1290
Trichogramma spp., 1290
Laricobius,
biological control, 896
competition, 896
predation, 896
larvae,
arrestment, 603
kairomone, 603
microencapsulated, 603
pear ester, 603
larval development,
dog-strangling vine, 841
fecundity, 841
host range, 841
Hypena, 841
larval developmental sites,
spatial variability, 213
Stomoxys calcitrans, 213
larval feeding,
larval movement, 1494
Striacosta albicosta, 1494
larval movement,
larval feeding, 1494
Striacosta albicosta, 1494
larval parasitoid,
Bactrocera oleae (Rossi), 497
irradiated host, 497
longevity, 497
parasitism, 497
Lasius alienus,
Aphaenogaster picea, 463
Aphaenogaster rudis, 463
biotic interactions, 463
community ecology, 463
leaf age,
Bemisia tabaci B biotype, 1125
host suitability, 1125
interspecific interaction, 1125
Trialeurodes vaporariorum, 1125
leaf traits,
indirect interactions, 1409
induced defenses, 1409

terpenes, 1409
vertebrate herbivory, 1409
Lepidoptera, 81, 849
biological control, 81
biology, 849
fertility life table, 849
host plant use, 81
invasive species, 81
oriental fruit moth, 849
Totricidae, 81
lesser cornstalk borer,
bioenergy, 865
ethanol, 865
herbivore, 865
pest management, 865
liberibacter,
Bactericera cockerelli, 1019
potato, 1019
psyllid haplotypes, 1019
zebra chip, 1019
life table, 30, 326, 1208, 1672
artificial diet, 30
beet armyworm, 326
Bemisia tabaci biotype B, 1672
Brontispa longissima, 1208
fecundity, 1208
growth and development, 1208
Helicoverpa armigera, 30
high-temperature stress, 1672
host plant, 1208
Iran, 326
long-term fitness effects, 1672
reproduction, 326
soybean, 326
Zea mays, 30
Lissorhoptrus oryzophilus,
antibiotics, 469
reproduction, 469
symbiont, 469
Wolbachia, 469
logistic,
phenology, 355
t-test, 355
Tyria jacobaeae, 355
long-term fitness effects,
Bemisia tabaci biotype B, 1672
high-temperature stress, 1672
life table, 1672
longevity, 20, 51, 497, 1062
Aonidiella aurantii, 51
Bactrocera oleae (Rossi), 497
Bailey's triple catch, 20
biological control, 51
bottom-up effects, 1062
cattle, 20
detritus, 1062
fluorescent marking, 20
forecasting, 20
irradiated host, 497
larval parasitoid, 497
mosquitoes, 1062
parasitism, 497
sex ratio, 51
size, 51
loss rate,
Bailey's triple-catch, 141
Jolly–Seber's stochastic method, 141
Monte-Carlo simulation, 141
population estimates, 141
lowbush blueberry,
ant diversity, 222
Formicidae, 222
Maine, 222
lower lethal temperature,
corn earworm, 172
lower lethal time, 172
supercooling point, 172
lower lethal time,
corn earworm, 172
lower lethal temperature, 172
supercooling point, 172
Loxostege sticticalis,
diapause, 1199
flight, 1199
reproduction, 1199
Lygodium microphyllum,
mating biology, 308
sexual selection, 308
weed biocontrol, 308
Lygus lineolaris, 1, 1302
age-specific life table, 1
development, 1
diapause, 1302
fecundity, 1
plant debris, 1302
tarnished plant bug, 1302
temperature, 1
winter host plants, 1302
Lygus,
cotton, 342
Gossypium, 342
western tarnished plant bug, 342
within-plant distribution, 342
lysozyme,
cannibalism, 166
disease, 166
katydid, 166
melanism, 166

M

macroinvertebrate, 802, 1322
biodiversity, 802
drainage ditch, 802
egg masses, 1322
hydrological seasonality, 1322
nutrients, 802
recruitment, 1322
tropical streams, 1322
water quality, 802
Macrosiphum euphorbiae,
microsatellites, 420
population structure, 420
year effect, 420
Macrotermes gilvus,
microsatellite, 426
next generation sequencing, 426
454 pyrosequencing, 426
termite, 426
Maine,
ant diversity, 222
Formicidae, 222
lowbush blueberry, 222
maize,
fall armyworm, 1131
Spodoptera frugiperda, 1131
sweet corn, 1131
Malaise trap,
Costa Rica, 1544
entomological net, 1544
immature stages, 1544
phytotelmata, 1544
mannitol,
aphid, 1638
overwintering, 1638
polyols, 1638
supercooling points, 1638
manuka oil,
electroantennography, 1597
phoebe oil, 1597
redbay ambrosia beetle, 1597
silkbay, 1597
mark–capture,
glassy-winged sharpshooter, 1279
insect dispersal, 1279
Pierce's disease, 1279
Xylella fastidiosa, 1279
massively parallel bacterial tag-encoded FLX-Titanium amplicon pyrosequencing,
microbiome, 98
psyllids, 98
pyrosequencing, 98
symbionts, 98
mate-location behavior,
Delphacidae, 1231
global warming, 1231
planthopper, 1231
temperature, 1231
mating biology,
Lygodium microphyllum, 308
sexual selection, 308
weed biocontrol, 308
mating disruption, 399, 407
apple, 399, 407
Cydia pomonella, 399, 407
monitoring, 399
plant volatiles, 407
sex pheromone, 399
traps, 407
mature larvae,
Bactrocera oleae, 1177
cultural control, 1177
fruit fly dispersal, 1177
teneral adult, 1177

mealybug,
biological control, 1184
parasitism, 1184
parasitoids, 1184
Mediterranean bioclimate,
abiotic factors, 785
dung beetle assemblages, 785
spatial scale, 785
temporal scale, 785
Megarhyssa,
Ibalia, 731
Pseudorhyssa, 731
Rhyssa, 731
Sirex noctilio, 731
melanism,
cannibalism, 166
disease, 166
katydid, 166
lysozyme, 166
Metarhizium anisopliae,
Burg El-Arab, 11
Egypt, 11
monthly abundance, 11
sand flies, 11
2-methyl-6-vinylpyrazine,
conspecifics, 387
host fruit, 387
papaya fruit fly, 387
pheromone, 387
microbiome,
massively parallel bacterial tag-encoded FLX-Titanium amplicon pyrosequencing, 98
psyllids, 98
pyrosequencing, 98
symbionts, 98
microencapsulated,
arrestment, 603
kairomone, 603
larvae, 603
pear ester, 603
microhabitat,
environmental stress, 979
geometric morphometrics, 979
preadaptation, 979
microhabitat preference,
aspen leaf miner, 180
overwintering, 180
Populus tremuloides, 180
supercooling, 180
microsatellite, 420, 426, 905
Bombus, 905
clover, 905
Macrosiphum euphorbiae, 420
Macrotermes gilvus, 426
next generation sequencing, 426
pollinators, 905
population structure, 420
454 pyrosequencing, 426
termite, 426
year effect, 420
mid-Atlantic agriculture,
ecosystem services, 813
native bees, 813
non-*Apis* bee crop pollination, 813
wild bees, 813
minute pirate bugs,
banker plants, 487
Frankliniella, 487
Orius, 487
thrips, 487
mirids,
CO-I, 1239
cryptic species, 1239
DNA barcoding, 1239
species-specific marker, 1239
monitoring, 399, 455, 1268
apple, 399
arthropod abundance, 1268
blueberry gall midge, 455
Bt corn, 1268
Cydia pomonella, 399
mating disruption, 399
phenology, 455
PRC analysis, 1268
riparian area, 1268
sex pheromone, 399
trap, 455
Monochamus,
Cerambycidae, 1587
host volatile, 1587
pheromone, 1587
2-(undecyloxy)-ethanol, 1587
Monte-Carlo simulation,
Bailey's triple-catch, 141
Jolly–Seber's stochastic method, 141
loss rate, 141
population estimates, 141
monthly abundance,
Burg El-Arab, 11
Egypt, 11
Metarhizium anisopliae, 11
sand flies, 11
mosquitoes,
bottom-up effects, 1062
Detritus, 1062
longevity, 1062
movement, 516, 578, 1426
Aphis glycines, 1426
dispersal, 578
foraging behavior, 516
harmonic radar, 578
host-plant resistance, 1426
Hylobius warreni, 578
partial refugia, 516
sampling, 578
tritrophic interactions, 516
within-plant distribution, 1426
mud-dauber wasps,
aggressive mimicry, 1474
anti-predator behavior, 1474
predator–prey interaction, 1474
Theridiidae, 1474
mulch,
Carabidae, 1097
compost, 1097
highbush blueberries, 1097
Staphylinidae, 1097
mutualism,
ant–plant interactions, 1405
elaiosome, 1405
myrmecochory, 1405
trophic interactions, 1405
mycetome,
barcode, 108
Periboeum pubescens, 108
symbiont, 108
Xylergates pulcher, 108
myrmecochory,
ant–plant interactions, 1405
elaiosome, 1405
mutualism, 1405
trophic interactions, 1405
myrmecophily,
aphid, 59
Homoptera, 59
invasive ant, 59
Myrmica rubra, 59
Myrmica rubra,
aphid, 59
Homoptera, 59
invasive ant, 59
myrmecophily, 59
Myzus ascalonicus,
biological invasion, 1398
host plant, 1398
Indian Oceanic Islands, 1398
Kerguelen Islands, 1398

N

native bees,
ecosystem services, 813
mid-Atlantic agriculture, 813
non-*Apis* bee crop pollination, 813
wild bees, 813
natural enemies, 289, 1364
apple, 1364
biological control, 1364
Deladenus siricidicola, 289
Eriosoma lanigerum, 1364
Ibalia leucospoides, 289
parasitic nematode, 289
parasitoids, 289
survey, 1364
natural enemy diversity,
aphids, 1077
flower diversity, 1077
natural pest suppression, 1077
natural pest suppression,
aphids, 1077
flower diversity, 1077
natural enemy diversity, 1077

natural selection,
body size, 1612
diapause, 1612
hibernation, 1612
polyandry, 1612
Neosilba,
Aganaspis, 233
Anastrepha, 233
Asobara, 233
Doryctobracon, 233
next generation sequencing,
Macrotermes gilvus, 426
microsatellite, 426
454 pyrosequencing, 426
termite, 426
niche overlap,
Ceutorhynchus alliariae, 1086
Ceutorhynchus roberti, 1086
competitive coexistence, 1086
niche partitioning, 1086
niche partitioning, 238, 1086
Ceutorhynchus alliariae, 1086
Ceutorhynchus roberti, 1086
competitive coexistence, 1086
exotic species, 238
feeding preference, 238
niche overlap, 1086
Onthophagus, 238
Phanaeus, 238
night collections,
Araneae, 1534
Atlantic Rainforest, 1534
day collections, 1534
sampling effort, 1534
nitric oxide,
entomopathogenic fungi, 959
immunity, 959
phenol oxidases, 959
scarabaeidae larvae, 959
nitrogen,
blue-stain, 478
Dendroctonus, 478
pine, 478
symbiosis, 478
Noctuidae,
biological control, 657
Braconidae, 657
Spodoptera exigua, 657
non-*Apis* bee crop pollination,
ecosystem services, 813
mid-Atlantic agriculture, 813
native bees, 813
wild bees, 813
nonbiting midge,
degree-day, 950
developmental rate, 950
thermal threshold, 950
toxicity test organism, 950
nonparametric multiplicative regression,
beet leafhopper, 768
Circulifer, 768
pest management, 768
purple top disease, 768
nontarget arthropods,
environmental risk assessment, 1687
genetically engineered plants, 1687
ladybird beetles, 1687
nontarget effects,
Cybocephalus spp., 377
Encarsia citrina, 377
euonymus scale (*Unaspis euonymi*), 377
nursery production, 377
nontarget risk assessment,
Bacillus thuringiensis, 1037
Cry3Bb1, 1037
Theridion impressum, 1037
nursery production,
Cybocephalus spp., 377
Encarsia citrina, 377
euonymus scale (*Unaspis euonymi*), 377
nontarget effects, 377
nutrients, 333, 802
alkaloids, 333
biodiversity, 802
coccid, 333
coffee plants, 333
drainage ditch, 802
macroinvertebrate, 802
phenolic compounds, 333
water quality, 802
Nylanderia pubens,
cDNA, 1008
ecotype, 1008
genomic DNA, 1008
Partrechina pubens, 1008

O

oilseed rape,
canola, 72
epigeic predators, 72
intercropping, 72
sustainable agriculture, 72
olfactometer, 125, 1169
cotton, 125
Harmonia axyridis, 1169
Hemiptera, 125
horsemint, 125
potassium metabisulfite, 1169
prior experience, 125
repellent, 1169
vineyard, 1169
olive fruit fly,
artificial diet, 265
invasion ecology, 265
sterile insect technology, 265
symbiosis, 265
Onthophagus,
exotic species, 238
feeding preference, 238
niche partitioning, 238
Phanaeus, 238
ontogenetic succession,
chemical defense, 541
deciduous forest, 541
herbivores, 541
insects, 541
oriental fruit moth,
biology, 849
fertility life table, 849
Lepidoptera, 849
orientation,
biocontrol, 1466
dispersal, 1466
Persicaria perfoliata, 1466
Rhinoncomimus latipes, 1466
Orius,
banker plants, 487
Frankliniella, 487
minute pirate bugs, 487
thrips, 487
Orius insidiosus,
Harmonia axyridis, 1565
integrated pest management, 1565
soybean aphid, 1565
spirotetramat, 1565
orthopteran,
cricket, 1644
feeding, 1644
Gryllus, 1644
reproduction, 1644
Oryza sativa,
Diatraea saccharalis, 571
host-plant resistance, 571
oviposition behavior, 571
Osmia rufa,
diapause, 1621
simulated climate change, 1621
temperature, 1621
Oulema melanopus, 761, 1516
degree-days, 761
integrated pest management, 761
predictive model, 761
SADIE, 1516
sampling, 761, 1516
spatial aggregation, 1516
spatial association, 1516
outbreaks,
ancestral state reconstruction, 723
bark beetles, 723
conifers, 723
Dendroctonus, 723
overwintering, 180, 1638
aphid, 1638
aspen leaf miner, 180
mannitol, 1638
microhabitat preference, 180
polyols, 1638
Populus tremuloides, 180

supercooling, 180
supercooling points, 1638
oviposition behavior,
Diatraea saccharalis, 571
host-plant resistance, 571
Oryza sativa, 571
oviposition preference,
insecticide residue, 880
Plutella xylostella, 880
resistance management, 880
transgenic plants, 880

P

pan trapping,
cucumber, 532
wild bees, 532
Wisconsin, 532
papaya fruit fly,
conspecifics, 387
host fruit, 387
2-methyl-6-vinylpyrazine, 387
pheromone, 387
paraquat,
common chickweed, 362
glyphosate, 362
Stellaria media, 362
tobacco thrips, 362
parasitic nematode,
Deladenus siricidicola, 289
Ibalia leucospoides, 289
natural enemies, 289
parasitoids, 289
parasitism, 497, 1184
Bactrocera oleae (Rossi), 497
biological control, 1184
irradiated host, 497
larval parasitoid, 497
longevity, 497
mealybug, 1184
parasitoids, 1184
parasitoids, 245, 289, 1184
Agrilus planipennis, 245
biological control, 1184
Deladenus siricidicola, 289
exotics and invasive species, 245
Ibalia leucospoides, 289
indigenous, 245
mealybug, 1184
natural enemies, 289
parasitic nematode, 289
parasitism, 1184
wood borers, 245
Parthenium hysterophorus,
biological control, 317
Carmenta ithacae, 317
CLIMEX model, 317
Sesiidae, 317
parthenogenesis,
development, 706
Echinothrips americanus, 706
sexual reproduction, 706
partial refugia,
foraging behavior, 516
movement, 516
tritrophic interactions, 516
Partrechina pubens,
cDNA, 1008
ecotype, 1008
genomic DNA, 1008
Nylanderia pubens, 1008
pear ester,
arrestment, 603
kairomone, 603
larvae, 603
microencapsulated, 603
pecan,
Beauveria bassiana, 298
biological control, 298
cover crop, 298
Curculio caryae, 298
Periboeum pubescens,
barcode, 108
mycetome, 108
symbiont, 108
Xylergates pulcher, 108
Persicaria perfoliata,
biocontrol, 1466
dispersal, 1466
orientation, 1466
Rhinoncomimus latipes, 1466
pest, 1029, 1505
agriculture, 1505
Diuraphis noxia, 1505
Fusarium, 1029
intraspecies mixture, 1029
predator, 1029
refuge, 1029
Russian wheat aphid, 1505
spatial model, 1505
pest management, 768, 865
beet leafhopper, 768
bioenergy, 865
Circulifer, 768
ethanol, 865
herbivore, 865
lesser cornstalk borer, 865
nonparametric multiplicative regression, 768
purple top disease, 768
pest resurgence,
conservation biological control, 621
Humulus lupulus, 621
Podosphaera macularis, 621
Phanaeus,
exotic species, 238
feeding preference, 238
niche partitioning, 238
Onthophagus, 238
phenol oxidases,
entomopathogenic fungi, 959
immunity, 959
nitric oxide, 959
scarabaeidae larvae, 959
phenolic compounds,
alkaloids, 333
coccid, 333
coffee plants, 333
nutrients, 333
phenology, 355, 448, 455, 1462
Apis mellifera, 1462
biological control, 448
blueberry gall midge, 455
climate match, 448
distribution, 1462
logistic, 355
monitoring, 455
Prokelisia marginata, 448
Spartina alterniflora, 448
swarm, 1462
trap, 455
t-test, 355
Tyria jacobaeae, 355
pheromone, 387, 562, 648, 887, 1481, 1587
aggregation, 887
Blissus insularis Barber, 887
Cerambycidae, 1587
Choristoneura rosaceana, 1481
conspecifics, 387
copulatory behavior, 562
courtship, 562
daily rhythm, 562
dose–response, 648
emerald ash borer, 648
exposure, 1481
Grapholita molesta, 1481
host fruit, 387
host-plant volatiles, 887
host volatile, 1587
lactone, 648
Monochamus, 1587
2-methyl-6-vinylpyrazine, 387
papaya fruit fly, 387
semiochemicals, 562
sexual behavior, 1481
semiochemicals, 562
2-(undecyloxy)-ethanol, 1587
Y-tube olfactometer, 887
(3Z)-dodecen-12-olide, 648
phoebe oil,
electroantennography, 1597
manuka oil, 1597
redbay ambrosia beetle, 1597
silkbay, 1597
phoretic,
beetle–fungi association, 255
Fusarium solani, 255
Scolytogenes birosimensis, 255
tree mortality, 255
photoperiod, 697, 1606
Adelphocoris suturalis, 1606
Apolygus lucorum, 1606
biological control, 697

diapause, 1606
ectoparasitoid, 697
Habrobracon hebetor, 697
reproductive diapause, 697
sensitive stage, 1606
phytotelmata,
Costa Rica, 1544
entomological net, 1544
immature stages, 1544
Malaise trap, 1544
Pierce's disease,
glassy-winged sharpshooter, 1279
insect dispersal, 1279
mark–capture, 1279
Xylella fastidiosa, 1279
Pieris,
resistance, 1153
Stephanitis, 1153
stomata, 1153
toughness, 1153
Pieris phillyreifolia,
P. japonica, 1145
resistance, 1145
S. pyrioides, 1145
Stephanitis takeyai, 1145
Pikonema alaskensis,
diel activity, 594
diurnal, 594
intracrown, 594
thermoregulation, 594
pine,
blue-stain, 478
Dendroctonus, 478
nitrogen, 478
symbiosis, 478
pine needles,
blueberry maggot, 370
compost, 370
fly emergence, 370
Rhagoletis mendax, 370
pine wood nematode,
adult trees, 828
environmental condition, 828
shading, 828
virulence, 828
Pinus albicaulis,
green leaf volatiles, 1575
Pinus contorta, 1575
semiochemicals, 1575
verbenone, 1575
Pinus,
funnel traps, 91
Sirex, 91
Tremex, 91
Urocerus, 91
Pinus contorta,
green leaf volatiles, 1575
Pinus albicaulis, 1575
semiochemicals, 1575
verbenone, 1575
P. japonica,
Pieris phillyreifolia, 1145
resistance, 1145
S. pyrioides, 1145
Stephanitis takeyai, 1145
plant bug,
development time, 415
diapause, 415
rearing, 415
plant debris,
diapause, 1302
Lygus lineolaris, 1302
tarnished plant bug, 1302
winter host plants, 1302
plant insect interactions,
bermudagrass, 1139
Eisenia fetida, 1139
feeding ecology, 1139
Scapteriscus, 1139
plant volatiles,
apple, 407
Cydia pomonella, 407
mating disruption, 407
traps, 407
planthopper,
Delphacidae, 1231
global warming, 1231
mate-location behavior, 1231
temperature, 1231
plant–insect interactions,
global change, 1435
jasmonic acid, 1435
salicylic acid, 1435
Plutella xylostella, 880, 1653
biological control, 1653
climate change, 1653
Diadegma insulare, 1653
fluctuating temperature, 1653
insecticide residue, 880
oviposition preference, 880
resistance management, 880
transgenic plants, 880
Poaceae,
conservation biological control, 275
energy sugarcane, 275
refuge, 275
stemborer, 275
Podosphaera macularis,
conservation biological control, 621
Humulus lupulus, 621
pest resurgence, 621
pollen loads,
exclusion experiments, 1163
female reproductive success, 1163
foraging behavior, 1163
pollinators,
Bombus, 905
clover, 905
microsatellites, 905
polyandry,
body size, 1612
diapause, 1612
hibernation, 1612
natural selection, 1612
polyols,
aphid, 1638
mannitol, 1638
overwintering, 1638
supercooling points, 1638
population dynamics, 1311, 1372, 1501
apparent mortality, 1311
Betula, 1372
cannibalism, 1501
flour conditioning, 1501
forest decline, 1372
host plant resistance, 1372
insect outbreaks, 1372
Quercus rubra, 1311
scar tissue, 1311
Tribolium castaneum, 1501
wood borers, 1311
population estimates,
Bailey's triple-catch, 141
Jolly–Seber's stochastic method, 141
loss rate, 141
Monte-Carlo simulation, 141
population genetics,
invasive species, 1680
Isoptera, 1680
Reticulitermes, 1680
social insect, 1680
population structure,
Macrosiphum euphorbiae, 420
microsatellites, 420
year effect, 420
Populus tremuloides,
aspen leaf miner, 180
microhabitat preference, 180
overwintering, 180
supercooling, 180
potassium metabisulfite,
Harmonia axyridis, 1169
olfactometer, 1169
repellent, 1169
vineyard, 1169
potato,
Bactericera cockerelli, 1019
liberibacter, 1019
psyllid haplotypes, 1019
zebra chip, 1019
PRC analysis,
arthropod abundance, 1268
Bt corn, 1268
monitoring, 1268
riparian area, 1268
preadaptation,
environmental stress, 979
geometric morphometrics, 979
microhabitat, 979

predation,
biological control, 896
competition, 896
Laricobius, 896
predation monitoring,
augmentation, 1527
biological control, 1527
ELISA, 1527
protein marking, 1527
predator, 669, 872, 1029
apple orchard, 872
aromatic plant, 872
arthropod community, 872
Bemisia tabaci, 669
biological control, 669
Delphastus catalinae, 669
diet, 669
Fusarium, 1029
herbivore, 872
intraspecies mixture, 1029
pest, 1029
refuge, 1029
predator–prey interaction,
aggressive mimicry, 1474
anti-predator behavior, 1474
mud-dauber wasps, 1474
Theridiidae, 1474
predictive model,
degree-days, 761
integrated pest management, 761
Oulema melanopus, 761
sampling, 761
prey density,
biological control, 508
foraging behavior, 508
habitat structure, 508
tritrophic interactions, 508
prior experience,
cotton, 125
Hemiptera, 125
horsemint, 125
olfactometer, 125
proboscis length,
Bombus, 1332
competition, 1332
corolla length, 1332
elevation, 1332
Prokelisia marginata,
biological control, 448
climate match, 448
phenology, 448
Spartina alterniflora, 448
protein marking,
augmentation, 1527
biological control, 1527
ELISA, 1527
predation monitoring, 1527
Pseudacteon,
agroforestry, 1107
Azteca instabilis, 1107
coffee, 1107
trait-meditated interactions, 1107
Pseudorhyssa,
Ibalia, 731
Megarhyssa, 731
Rhyssa, 731
Sirex noctilio, 731
psyllids,
massively parallel bacterial tag-encoded FLX-Titanium amplicon pyrosequencing, 98
microbiome, 98
pyrosequencing, 98
symbionts, 98
psyllid haplotypes,
Bactericera cockerelli, 1019
liberibacter, 1019
potato, 1019
zebra chip, 1019
purple top disease,
beet leafhopper, 768
Circulifer, 768
nonparametric multiplicative regression, 768
pest management, 768
pyrosequencing,
massively parallel bacterial tag-encoded FLX-Titanium amplicon pyrosequencing, 98
microbiome, 98
psyllids, 98
symbionts, 98
454 pyrosequencing,
Macrotermes gilvus, 426
microsatellite, 426
next generation sequencing, 426
termite, 426

Q

Quercus rubra,
apparent mortality, 1311
population dynamics, 1311
scar tissue, 1311
wood borers, 1311

R

rabid wolf spider,
thermal ecology, 1631
thermal performance, 1631
real-time qPCR,
bacterial community, 1386
Brevicoryne brassicae, 1386
endosymbiont, 1386
16S ribosomal DNA, 1386
rearing,
development time, 415
diapause, 415
plant bug, 415
recruitment,
egg masses, 1322
hydrological seasonality, 1322
macroinvertebrates, 1322
tropical streams, 1322
redbay ambrosia beetle,
electroantennography, 1597
manuka oil, 1597
phoebe oil, 1597
silkbay, 1597
refuge, 275, 1029
conservation biological control, 275
energy sugarcane, 275
Fusarium, 1029
intraspecies mixture, 1029
pest, 1029
Poaceae, 275
predator, 1029
stemborer, 275
relative humidity,
activity, 587
Braconidae, 587
Ichneumonidae, 587
temperature, 587
repellent, 855, 1169
bumble bee, 855
floral volatile, 855
greenhouse, 855
Harmonia axyridis, 1169
olfactometer, 1169
potassium metabisulfite, 1169
tomato, 855
vineyard, 1169
reproduction, 326, 469, 989, 1199, 1644
antibiotics, 469
beet armyworm, 326
biological control, 989
cricket, 1644
development, 989
diapause, 1199
feeding, 1644
flight, 1199
Gryllus, 1644
Iran, 326
life table, 326
Lissorhoptrus oryzophilus, 469
Loxostege sticticalis, 1199
orthopteran, 1644
southern Africa, 989
soybean, 326
symbiont, 469
temperature, 989
Wolbachia, 469
reproductive development,
temperature, 941
western tarnished plant bug, 941
reproductive diapause,
biological control, 697
ectoparasitoid, 697

Habrobracon hebetor, 697
photoperiod, 697
resistance, 1145, 1153
Pieris phillyreifolia, 1145
P. japonica, 1145
Pieris, 1153
S. pyrioides, 1145
Stephanitis, 1153
Stephanitis takeyai, 1145
stomata, 1153
toughness, 1153
resistance management, 200, 880, 1255
Bt corn, 200
Bt cowpea, 1255
insecticide residue, 880
oviposition preference, 880
Plutella xylostella, 880
simulation, 200, 1255
transgenic plants, 880
Reticulitermes,
invasive species, 1680
Isoptera, 1680
population genetics, 1680
social insect, 1680
Rhagoletis mendax,
blueberry maggot, 370
compost, 370
fly emergence, 370
pine needles, 370
Rhinoncomimus latipes,
biocontrol, 1466
dispersal, 1466
orientation, 1466
Persicaria perfoliata, 1466
Rhyssa,
Ibalia, 731
Megarhyssa, 731
Pseudorhyssa, 731
Sirex noctilio, 731
riparian area,
arthropod abundance, 1268
Bt corn, 1268
monitoring, 1268
PRC analysis, 1268
root-feeding insect,
Delia radicum, 159
soil moisture, 159
soil temperature, 159
Russian wheat aphid,
agriculture, 1505
Diuraphis noxia, 1505
pest, 1505
spatial model, 1505

S

16S ribosomal DNA,
bacterial community, 1386
Brevicoryne brassicae, 1386
endosymbiont, 1386
real-time qPCR, 1386
SADIE,
Oulema melanopus, 1516
sampling, 1516
spatial aggregation, 1516
spatial association, 1516
salicylic acid, 1435, 1443
Bemisia tabaci, 1443
global change, 1435
jasmonic acid, 1435, 1443
plant–insect interactions, 1435
tomato, 1443
whitefly preconditioning, 1443
sampling, 578, 761, 1516
degree-days, 761
dispersal, 578
harmonic radar, 578
Hylobius warreni, 578
integrated pest management, 761
movement, 578
Oulema melanopus, 761, 1516
predictive model, 761
SADIE, 1516
spatial aggregation, 1516
spatial association, 1516
sampling effort,
Araneae, 1534
Atlantic Rainforest, 1534
day collections, 1534
night collections, 1534
sand flies,
Burg El-Arab, 11
Egypt, 11
Metarhizium anisopliae, 11
monthly abundance, 11
saproxylic,
Cerambycidae, 1350
host finding, 1350
Ips, 1350
Scolytinae, 1350
Scapteriscus,
bermudagrass, 1139
Eisenia fetida, 1139
feeding ecology, 1139
plant insect interactions, 1139
scar tissue,
apparent mortality, 1311
population dynamics, 1311
Quercus rubra, 1311
wood borers, 1311
scarabaeidae larvae,
entomopathogenic fungi, 959
immunity, 959
nitric oxide, 959
phenol oxidases, 959
Scolytinae, 392, 1350
Cerambycidae, 1350
Coleoptera, 392
Curculionidae, 392
host finding, 1350
Ips, 1350
saproxylic, 1350
secondary metabolites, 392
terpenes, 392
Scolytogenes birosimensis,
beetle–fungi association, 255
Fusarium solani, 255
phoretic, 255
tree mortality, 255
Scymnus camptodromus,
egg diapause, 1662
hatch, 1662
temperature, 1662
seasonal ecology,
Lambdina fiscellaria, 1290
thermal requirement, 1290
thermal response, 1290
Trichogramma spp., 1290
seasonality,
soil, 916
springtail, 916
walking, 916
second-generation sequencing,
single nucleotide polymorphism, 997
Striacosta albicosta, 997
transcriptome, 997
western bean cutworm, 997
secondary metabolites,
Coleoptera, 392
Curculionidae, 392
Scolytinae, 392
terpenes, 392
semiochemicals, 133, 562, 1575
ambrosia beetles, 133
copulatory behavior, 562
courtship, 562
daily rhythm, 562
green leaf volatiles, 1575
pheromones, 562
Pinus albicaulis, 1575
Pinus contorta, 1575
traps, 133
verbenone, 1575
sensitive stage,
Adelphocoris suturalis, 1606
Apolygus lucorum, 1606
diapause, 1606
photoperiod, 1606
Sesiidae,
biological control, 317
Carmenta ithacae, 317
CLIMEX model, 317
Parthenium hysterophorus, 317
sex pheromone,
apple, 399
Cydia pomonella, 399
mating disruption, 399
monitoring, 399
sex ratio,
Aonidiella aurantii, 51
biological control, 51
longevity, 51
size, 51

sexual behavior,
Choristoneura rosaceana, 1481
exposure, 1481
Grapholita molesta, 1481
pheromone, 1481
sexual reproduction,
development, 706
Echinothrips americanus, 706
parthenogenesis, 706
sexual selection,
Lygodium microphyllum, 308
mating biology, 308
weed biocontrol, 308
shading,
adult trees, 828
environmental condition, 828
pine wood nematode, 828
virulence, 828
silkbay,
electroantennography, 1597
manuka oil, 1597
phoebe oil, 1597
redbay ambrosia beetle, 1597
simulated climate change,
diapause, 1621
Osmia rufa, 1621
temperature, 1621
simulation, 200, 1255
Bt corn, 200
Bt cowpea, 1255
resistance management, 200, 1255
single nucleotide polymorphism,
second-generation sequencing, 997
Striacosta albicosta, 997
transcriptome, 997
western bean cutworm, 997
Sirex,
funnel traps, 91
Pinus, 91
Tremex, 91
Urocerus, 91
Sirex noctilio,
Ibalia, 731
Megarhyssa, 731
Pseudorhyssa, 731
Rhyssa, 731
size,
Aonidiella aurantii, 51
biological control, 51
longevity, 51
sex ratio, 51
small hive beetle larvae,
Aethina tumida, 152
honey bees, 152
insect trapping, 152
social insect,
invasive species, 1680
Isoptera, 1680
population genetics, 1680
Reticulitermes, 1680
Sogatella furcifera,
Cardinium, 833
cytoplasmic incompatibility, 833
infection frequency, 833
Wolbachia, 833
soil,
seasonality, 916
springtail, 916
walking, 916
soil moisture,
Delia radicum, 159
root-feeding insect, 159
soil temperature, 159
soil pH,
entomopathogen, 1115
Entomophaga maimaiga, 1115
fungal epizootic, 1115
gypsy moth, 1115
soil temperature,
Delia radicum, 159
root-feeding insect, 159
soil moisture, 159
soil-dwelling pest,
autecology, 966
habitat, 966
integrated pest management (IPM), 966
solid phase microextraction, 636, 933
ambrosia beetles, 636
chestnut volatile organic compound profile, 933
Curculio, 933
electroantennogram, 933
ethanol, 636
gas chromatography-mass spectrometry, 933
Xylosandrus germanus, 636
southern Africa,
biological control, 989
development, 989
reproduction, 989
temperature, 989
soybean, 282, 326, 750, 1417
Aphis glycines, 282, 750, 1417
beet armyworm, 326
Binodoxys communis, 282, 1417
biological control, 282, 1417
Cerotoma trifurcata, 750
Empoasca fabae, 750
host-plant resistance, 282, 1417
Iran, 326
life table, 326
reproduction, 326
winter rye, 750
soybean aphid,
Harmonia axyridis, 1565
integrated pest management, 1565
Orius insidiosus, 1565
spirotetramat, 1565
S. pyrioides,
Pieris phillyreifolia, 1145
P. japonica, 1145
resistance, 1145
Stephanitis takeyai, 1145
Spartina alterniflora,
biological control, 448
climate match, 448
phenology, 448
Prokelisia marginata, 448
Spathius galinae,
Agrilus planipennis, 925
biological control, 925
interspecific competition, 925
Tetrastichus planipennisi, 925
spatial aggregation,
Oulema melanopus, 1516
SADIE, 1516
sampling, 1516
spatial association, 1516
spatial association,
Oulema melanopus, 1516
SADIE, 1516
sampling, 1516
spatial aggregation, 1516
spatial model,
agriculture, 1505
Diuraphis noxia, 1505
pest, 1505
Russian wheat aphid, 1505
spatial pattern,
dispersal, 1050
geostatistics, 1050
Gomphocerinae, 1050
Italian locust, 1050
spatial scale,
abiotic factors, 785
dung beetle assemblages, 785
Mediterranean bioclimate, 785
temporal scale, 785
spatial variability,
larval developmental sites, 213
Stomoxys calcitrans, 213
species-specific marker,
CO-I, 1239
cryptic species, 1239
DNA barcoding, 1239
mirids, 1239
spirotetramat,
Harmonia axyridis, 1565
integrated pest management, 1565
Orius insidiosus, 1565
soybean aphid, 1565
Spodoptera exigua,
biological control, 657
Braconidae, 657
Noctuidae, 657
Spodoptera frugiperda,
fall armyworm, 1131
maize, 1131
sweet corn, 1131

springtail,
seasonality, 916
soil, 916
walking, 916
SSI model,
body mass, 714
development rate, 714
fitness, 714
intrinsic rate of increase, 714
Staphylinidae,
Carabidae, 1097
compost, 1097
highbush blueberries, 1097
mulch, 1097
Stellaria media,
common chickweed, 362
glyphosate, 362
paraquat, 362
tobacco thrips, 362
stemborer,
conservation biological control, 275
energy sugarcane, 275
Poaceae, 275
refuge, 275
Stephanitis takeyai,
Pieris phillyreifolia, 1145
P. japonica, 1145
resistance, 1145
S. pyrioides, 1145
Stephanitis,
Pieris, 1153
resistance, 1153
stomata, 1153
toughness, 1153
sterile insect technology,
artificial diet, 265
invasion ecology, 265
olive fruit fly, 265
symbiosis, 265
stomata,
Pieris, 1153
resistance, 1153
Stephanitis, 1153
toughness, 1153
Stomoxys calcitrans,
larval developmental sites, 213
spatial variability, 213
Striacosta albicosta, 997, 1494
larval feeding, 1494
larval movement, 1494
second-generation sequencing, 997
single nucleotide polymorphism, 997
transcriptome, 997
western bean cutworm, 997
supercooling,
aspen leaf miner, 180
microhabitat preference, 180
overwintering, 180
Populus tremuloides, 180
supercooling point,
corn earworm, 172
lower lethal temperature, 172
lower lethal time, 172
supercooling points,
aphid, 1638
mannitol, 1638
overwintering, 1638
polyols, 1638
survey,
apple, 1364
biological control, 1364
Eriosoma lanigerum, 1364
natural enemies, 1364
survival rate,
Carposina sasakii, 349
duration, 349
fecundity, 349
host plant, 349
sustainable agriculture,
canola, 72
epigeic predators, 72
intercropping, 72
oilseed rape, 72
swallow-wort,
Aulacorthum solani, 665
foxglove aphid, 665
host suitability, 665
Vincetoxicum, 665
swarm,
Apis mellifera, 1462
distribution, 1462
phenology, 1462
sweet corn,
fall armyworm, 1131
maize, 1131
Spodoptera frugiperda, 1131
symbiont, 98, 108, 469
antibiotics, 469
barcode, 108
Lissorhoptrus oryzophilus, 469
massively parallel bacterial tag-encoded FLX-Titanium amplicon pyrosequencing, 98
microbiome, 98
mycetome, 108
Peribocum pubescens, 108
psyllids, 98
pyrosequencing, 98
reproduction, 469
Wolbachia, 469
Xylergates pulcher, 108
symbiosis, 265, 478
artificial diet, 265
blue-stain, 478
Dendroctonus, 478
invasion ecology, 265
nitrogen, 478
olive fruit fly, 265
pine, 478
sterile insect technology, 265
sympatric speciation,
Asphondylia borrichiae, 1246
host-associated divergence, 1246
host-race formation, 1246

T

tarnished plant bug,
diapause, 1302
Lygus lineolaris, 1302
plant debris, 1302
winter host plants, 1302
temperature, 1, 587, 676, 941, 989, 1190, 1215, 1231, 1621, 1662
activity, 587
age-specific life table, 1
Bactericera cockerelli, 1190
biological control, 989
Braconidae, 587
Coccinella septempunctata, 676
Delphacidae, 1231
development, 1, 989, 1190
diapause, 1621
egg diapause, 1662
electrical penetration graph, 1215
fat body, 676
fecundity, 1
global warming, 1231
Harmonia axyridis, 676
hatch, 1662
Homalodisca vitripennis, 1215
Ichneumonidae, 587
ingestion, 1215
insect feeding, 1215
Lygus lineolaris, 1
mate-location behavior, 1231
Osmia rufa, 1621
planthopper, 1231
relative humidity, 587
reproduction, 989
reproductive development, 941
Scymnus camptodromus, 1662
simulated climate change, 1621
southern Africa, 989
tomato psyllid (potato psyllid), 1190
weight, 676
western tarnished plant bug, 941
temporal scale,
abiotic factors, 785
dung beetle assemblages, 785
Mediterranean bioclimate, 785
spatial scale, 785
teneral adult,
Bactrocera oleae, 1177
cultural control, 1177
fruit fly dispersal, 1177
mature larvae, 1177

termite,
Macrotermes gilvus, 426
microsatellite, 426
next generation sequencing, 426
454 pyrosequencing, 426
terpenes, 392, 1409
Coleoptera, 392
Curculionidae, 392
indirect interactions, 1409
induced defenses, 1409
leaf traits, 1409
Scolytinae, 392
secondary metabolites, 392
vertebrate herbivory, 1409
Tetrastichus planipennisi,
Agrilus planipennis, 925
biological control, 925
interspecific competition, 925
Spathius galinae, 925
Theridiidae,
aggressive mimicry, 1474
anti-predator behavior, 1474
mud-dauber wasps, 1474
predator–prey interaction, 1474
Theridion impressum,
Bacillus thuringiensis, 1037
Cry3Bb1, 1037
nontarget risk assessment, 1037
thermal ecology,
rabid wolf spider, 1631
thermal performance, 1631
thermal performance,
rabid wolf spider, 1631
thermal ecology, 1631
thermal requirement,
Lambdina fiscellaria, 1290
seasonal ecology, 1290
thermal response, 1290
Trichogramma spp., 1290
thermal response,
Lambdina fiscellaria, 1290
seasonal ecology, 1290
thermal requirement, 1290
Trichogramma spp., 1290
thermal threshold,
degree-day, 950
developmental rate, 950
nonbiting midge, 950
toxicity test organism, 950
thermoregulation,
diel activity, 594
diurnal, 594
intracrown, 594
Pikonema alaskensis, 594
thrips,
banker plants, 487
Frankliniella, 487
minute pirate bugs, 487
Orius, 487
tobacco thrips,
common chickweed, 362
glyphosate, 362
paraquat, 362
Stellaria media, 362
tomato, 855, 1443
Bemisia tabaci, 1443
bumble bee, 855
floral volatile, 855
greenhouse, 855
jasmonic acid, 1443
repellent, 855
salicylic acid, 1443
whitefly preconditioning, 1443
tomato psyllid (potato psyllid),
Bactericera cockerelli, 1190
development, 1190
temperature, 1190
Totricidae,
biological control, 81
host plant use, 81
invasive species, 81
Lepidoptera, 81
toughness,
Pieris, 1153
resistance, 1153
Stephanitis, 1153
stomata, 1153
toxicity test organism,
degree-day, 950
developmental rate, 950
nonbiting midge, 950
thermal threshold, 950
trait-meditated interactions,
agroforestry, 1107
Azteca instabilis, 1107
coffee, 1107
Pseudacteon, 1107
transcriptome,
second-generation sequencing, 997
single nucleotide polymorphism, 997
Striacosta albicosta, 997
western bean cutworm, 997
transgenic plants,
insecticide residue, 880
oviposition preference, 880
Plutella xylostella, 880
resistance management, 880
trap, 133, 407, 455
ambrosia beetles, 133
apple, 407
blueberry gall midge, 455
Cydia pomonella, 407
mating disruption, 407
monitoring, 455
phenology, 455
plant volatiles, 407
semiochemicals, 133
tree mortality,
beetle–fungi association, 255
Fusarium solani, 255
phoretic, 255
Scolytogenes birosimensis, 255
Tremex,
funnel traps, 91
Pinus, 91
Sirex, 91
Urocerus, 91
Trialeurodes vaporariorum,
Bemisia tabaci B biotype, 1125
host suitability, 1125
interspecific interaction, 1125
leaf age, 1125
Tribolium castaneum, 188, 1501
bottleneck, 188
cannibalism, 1501
flour conditioning, 1501
F_{ST}, 188
genetic fingerprinting, 188
genetic structure, 188
population dynamics, 1501
Trichogramma spp.,
Lambdina fiscellaria, 1290
seasonal ecology, 1290
thermal requirement, 1290
thermal response, 1290
tritrophic interactions, 508, 516
biological control, 508
foraging behavior, 508, 516
habitat structure, 508
movement, 516
partial refugia, 516
prey density, 508
trophic interactions,
ant–plant interactions, 1405
elaiosome, 1405
mutualism, 1405
myrmecochory, 1405
tropical streams,
egg masses, 1322
hydrological seasonality, 1322
macroinvertebrates, 1322
recruitment, 1322
Tsuga canadensis,
Adelges tsugae, 523
false rings, 523
Fiorinia externa, 523
herbivory, 523
t-test,
logistic, 355
phenology, 355
Tyria jacobaeae, 355
Tyria jacobaeae,
logistic, 355
phenology, 355
t-test, 355

U

2-(undecyloxy)-ethanol,
Cerambycidae, 1587
host volatile, 1587
Monochamus, 1587
pheromone, 1587

Urocerus,
funnel traps, 91
Pinus, 91
Sirex, 91
Tremex, 91

V

vegetation,
Black Hills, 1069
Carabidae, 1069
Coleoptera, 1069
ground beetles, 1069
verbenone,
green leaf volatiles, 1575
Pinus albicaulis, 1575
Pinus contorta, 1575
semiochemicals, 1575
vertebrate herbivory,
indirect interactions, 1409
induced defenses, 1409
leaf traits, 1409
terpenes, 1409
Vincetoxicum,
Aulacorthum solani, 665
foxglove aphid, 665
host suitability, 665
swallow-wort, 665
vineyard,
Harmonia axyridis, 1169
olfactometer, 1169
potassium metabisulfite, 1169
repellent, 1169
virulence,
adult trees, 828
environmental condition, 828
pine wood nematode, 828
shading, 828

W

walking,
seasonality, 916
soil, 916
springtail, 916
waste management,
adult emergence, 971
egg clustering, 971
egg desiccation hypothesis, 971
egg eclosion, 971
water quality,
biodiversity, 802
drainage ditch, 802
macroinvertebrate, 802
nutrients, 802
weed biocontrol,
Lygodium microphyllum, 308
mating biology, 308
sexual selection, 308
weight,
Coccinella septempunctata, 676
fat body, 676
Harmonia axyridis, 676
temperature, 676
western bean cutworm,
second-generation sequencing, 997
single nucleotide polymorphism, 997
Striacosta albicosta, 997
transcriptome, 997
western tarnished plant bug, 342, 941
cotton, 342
Gossypium, 342
Lygus, 342
reproductive development, 941
temperature, 941
within-plant distribution, 342
whitefly preconditioning,
Bemisia tabaci, 1443
jasmonic acid, 1443
salicylic acid, 1443
tomato, 1443
wild bees, 532, 813
cucumber, 532
ecosystem services, 813
mid-Atlantic agriculture, 813
native bees, 813
non-*Apis* bee crop pollination, 813
pan trapping, 532
Wisconsin, 532
winter host plants,
diapause, 1302
Lygus lineolaris, 1302
plant debris, 1302
tarnished plant bug, 1302
winter rye,
Aphis glycines, 750
Cerotoma trifurcata, 750
Empoasca fabae, 750
soybean, 750
Wisconsin,
cucumber, 532
pan trapping, 532
wild bees, 532
within-plant distribution, 342, 1426
Aphis glycines, 1426
cotton, 342
Gossypium, 342
host-plant resistance, 1426
Lygus, 342
movement, 1426
western tarnished plant bug, 342
Wolbachia, 469, 833
antibiotics, 469
Cardinium, 833
cytoplasmic incompatibility, 833
infection frequency, 833
Lissorhoptrus oryzophilus, 469
reproduction, 469
Sogatella furcifera, 833
symbiont, 469
wood borers, 245, 1311
Agrilus planipennis, 245
apparent mortality, 1311
exotics and invasive species, 245
indigenous, 245
parasitoids, 245
population dynamics, 1311
Quercus rubra, 1311
scar tissue, 1311

X

Xyleborini ambrosia beetles,
agricultural areas, 776
diversity, 776
flight pattern, 776
Xylella fastidiosa,
glassy-winged sharpshooter, 1279
insect dispersal, 1279
mark–capture, 1279
Pierce's disease, 1279
Xylergates pulcher,
barcode, 108
mycetome, 108
Periboeum pubescens, 108
symbiont, 108
Xylosandrus germanus,
ambrosia beetles, 636
ethanol, 636
solid phase microextraction, 636

Y

year effect,
Macrosiphum euphorbiae, 420
microsatellites, 420
population structure, 420
Y-tube olfactometer,
aggregation, 887
Blissus insularis Barber, 887
host-plant volatiles, 887
pheromones, 887

Z

Zea mays,
artificial diet, 30
Helicoverpa armigera, 30
life table, 30
zebra chip,
Bactericera cockerelli, 1019
liberibacter, 1019
potato, 1019
psyllid haplotypes, 1019
(3*Z*)-dodecen-12-olide,
dose–response, 648
emerald ash borer, 648
lactone, 648
pheromone, 648